

VU Research Portal

Governing processes for reactive nitrogen compounds in the atmosphere in relation to ecosystem climatic and human health impacts

Hertel, O.; Skjøth, C.A.; Reis, S.; Bleeker, A.; Harrison, R.; Cape, J.N.; Fowler, D.; Skiba, U.; Simpson, D.; Jickells, T.; Kulmala, M.; Gyldenærne, S.; Sørensen, L.L.; Erisman, J.W.; Sutton, M.A.

published in

Biogeosciences Discussions
2012

DOI (link to publisher)

[10.5194/bgd-9-9349-2012](https://doi.org/10.5194/bgd-9-9349-2012)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Hertel, O., Skjøth, C. A., Reis, S., Bleeker, A., Harrison, R., Cape, J. N., Fowler, D., Skiba, U., Simpson, D., Jickells, T., Kulmala, M., Gyldenærne, S., Sørensen, L. L., Erisman, J. W., & Sutton, M. A. (2012). Governing processes for reactive nitrogen compounds in the atmosphere in relation to ecosystem climatic and human health impacts. *Biogeosciences Discussions*, 9(7), 9349-9423. <https://doi.org/10.5194/bgd-9-9349-2012>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Governing processes for reactive nitrogen compounds in the atmosphere in relation to ecosystem, climatic and human health impacts

**O. Hertel^{1,2}, C. A. Skjøth^{1,3}, S. Reis⁴, A. Bleeker⁵, R. Harrison^{6,7}, J. N. Cape⁴,
D. Fowler⁴, U. Skiba⁴, D. Simpson^{8,9}, T. Jickells¹⁰, M. Kulmala¹¹,
S. Gyldenkerne¹, L. L. Sørensen¹, J. W. Erisman¹², and M. A. Sutton⁴**

¹Department of Environmental Science, Aarhus University, P.O. Box 358,
Frederiksborgvej 399, 4000 Roskilde, Denmark

²Department for Environmental, Social and Spatial Change (ENSPAC), Roskilde University,
4000 Roskilde, Denmark

³Department of Physical Geography and Ecosystems Science, Lund University,
Sölvegatan 12, 223 62 Lund, Sweden

⁴Centre for Ecology & Hydrology (CEH), Bush Estate Penicuik Midlothian EH26 0QB, UK

⁵Energy Research Centre of the Netherlands (ECN), Biomass, Coal & Environmental
Research, P.O. Box 19, 1755 ZG Petten, The Netherlands

⁶School of Geography, Earth and Environmental Sciences, University of Birmingham,
Edgbaston Birmingham B15 2TT, UK

⁷Department of Environmental Sciences/Center of Excellence in Environmental Studies, King Abdulaziz University, P.O. Box 80203, Jeddah, 21589, Saudi Arabia

⁸Norwegian Meteorological Institute, Henrik Mons Plass 1, 0313 Oslo, Norway

⁹Department of Earth & Space Sciences, Chalmers University of Technology, 412 96 Gothenburg, Sweden

¹⁰University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK

¹¹Department of Physics, University of Helsinki, Finland

¹²Earth and Climate, VU University Amsterdam and Louis Bolk Institute, Hoofdweg 24, 3972 LA Driebergen, The Netherlands

Received: 15 July 2012 – Accepted: 16 July 2012 – Published: 27 July 2012

Correspondence to: O. Hertel (ole.hertel@dmu.dk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Reactive nitrogen (N_r) compounds have different fates in the atmosphere due to differences in governing processes of physical transport, deposition and chemical transformation. N_r compounds addressed here include reduced nitrogen (NH_x : ammonia (NH_3) and its reaction product ammonium (NH_4^+)), oxidized nitrogen (NO_y : nitrogen monoxide (NO) + nitrogen dioxide (NO_2) and their reaction products) as well as organic nitrogen compounds (organic N). Pollution abatement strategies need to take into account these differences in the governing processes of these compounds when assessing their impact on ecosystem services, biodiversity, human health and climate.

NO_x (NO + NO_2) emitted from traffic affects human health in urban areas where the presence of buildings increases the residence time in streets. In urban areas this leads to enhanced exposure of the population to NO_x concentrations. NO_x emissions have little impact on nearby ecosystems because of the small dry deposition rates of NO_x . These compounds need to be converted into nitric acid (HNO_3) before removal through deposition is efficient. HNO_3 sticks quickly to any surface and is thereby either dry deposited or incorporated into aerosols as nitrate (NO_3^-). In contrast to NO_x compounds, NH_3 has potentially high impacts on ecosystems near the main agricultural sources of NH_3 because of its large ground-level concentrations along with large dry deposition rates. Aerosol phase NH_4^+ and NO_3^- contribute significantly to background $PM_{2.5}$ and PM_{10} (mass of aerosols with a diameter of less than 2.5 and 10 μm , respectively) with an impact on radiation balance as well as potentially on human health. Little is known quantitatively and qualitatively about organic N in the atmosphere, other than that it contributes a significant fraction of wet-deposited N, and is present in both gaseous and particulate forms in the atmosphere. Further studies are needed to characterize the sources, air chemistry and removal rates of organic N emissions.

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1 Introduction

Reactive nitrogen compounds (N_r) affect ecosystem health (Sutton et al., 2011), human health (Aneja et al., 2009) and contribute to climate change (Butterbach-Bah et al., 2011; Erisman et al., 2011; Xu and Penner, 2012). However, the actual impact of specific emissions depends on a cascade of competing processes taking place simultaneously, which transforms pollutants along multiple paths, with multiple impacts (Galloway et al., 2003). One example is the release of NH_3 . Strong NH_3 sources contribute to high N loads to nearby ecosystems through a fast dry deposition rate to nearby surfaces (Hertel et al., 2006). This process competes with reactions that lead to the formation of aerosol phase NH_4^+ . The latter contributes to $PM_{2.5}$ and PM_{10} (mass of aerosols with an aerodynamic diameter less than 2.5 and 10 μm , respectively) with much longer transport distance and with potential health effects and impact on the radiation balance. The NH_4^+ -containing aerosols have low dry deposition velocities, can be transported up to 1000 km away and are mainly wet scavenged. Figure 1 provides an overview of the N_r pathways, with an emphasis on atmospheric N deposition. NH_3 is emitted mainly from agricultural sources (Bouwman et al., 1997), and has a typical atmospheric residence time of 24 h. Both NH_3 and gaseous NO_x lead to formation of aerosol phase compounds (NH_4^+ and NO_3^- , respectively) which have atmospheric residence times of several days. Furthermore NH_3 can take part in bi-directional surface fluxes (Sutton et al., 1995); in areas with large nitrogen (N) inputs, NH_3 may be released from plants and soil, and in areas with high gas-phase concentrations, deposition rates may be limited by surface saturation, and re-emission of absorbed NH_3 may occur (Flechard et al., 1999; Massad et al., 2010b; Nemitz et al., 2004a; Sutton et al., 2009a, b). By comparison, the primary form in which oxidized nitrogen (NO_y) is emitted is nitrogen monoxide (NO), which reacts to form nitrogen dioxide (NO_2) over minutes and hours depending on local oxidant chemistry (interactions are described well in various text books see e.g. Seinfeld and Pandis, 2006). Both of these compounds (especially NO) are removed slowly by dry deposition to vegetation. The main

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

scavenging process is therefore by chemical conversion. In the case of NO_2 , chemical reaction with hydroxyl (OH) radical forms nitric acid (HNO_3) with a typical transformation rate of 5 % per hour (Hertel, 1995). HNO_3 has a short atmospheric lifetime (on the order of seconds to hours), since it is quickly scavenged by uptake in aerosols, reaction with NH_3 or by dry deposition (it typically sticks to any surface – aerosol as well as vegetation and the ground) (Hertel et al., 2006). A US study (Day et al., 2008) found a lifetime of HNO_3 of 2.5 h for a mountain area in a sunny and warm climate, but the lifetime depends strongly on the environment and another study has shown lifetimes of 7–20 min in air masses processed by clouds containing ice-crystals (von Kuhlmann and Lawrence, 2006). Uptake of HNO_3 into existing aerosols or formation of new aerosols by reaction with NH_3 leads to aerosol bound nitrate (NO_3^-) as does the reaction with sea-salt (Pryor and Sorensen, 2000). Such ammonium (NH_4^+) and NO_3^- containing aerosols are mainly scavenged by wet deposition, due to the relatively small dry deposition rate of submicron aerosols (Gallagher et al., 2002). Aerosol phase NH_4^+ and NO_3^- may under certain circumstances – depending on humidity and temperature – be released back to gas phase NH_3 and HNO_3 . The greenhouse gas nitrous oxide (N_2O) is excluded in the figure, as it has little impact on atmospheric N deposition. To maintain simplicity of the diagram, reservoir compounds (e.g. HONO , HO_2NO_2 , PAN and PAN-like species) are also disregarded in Fig. 1, although these are discussed in the subsequent sections. The actual split between various pathways depends on the precise location where local climate conditions, local emission densities and distance from high emission areas all determine the overall fate of N_r . Another example of atmospheric processes affecting the impact of N_r is the complex flow conditions in urban street canyons. The generally short residence time of air in urban streets means that only fast reactions of NO_x are relevant. However, the presence of buildings, especially in streets with close building facades on both sides of the street, increases the local residence time of traffic pollutants emitted in the street sufficiently to significantly increase also the exposure of the population (Hertel and Goodsites, 2009). These processes – the flow conditions and the chemical conversion – affect the ratio between NO

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



which has little direct impact on human health and the airway irritant NO₂ that e.g. may enhance impact on airborne allergens (Tunncliffe et al., 1994). As can be seen from this short introduction, the atmospheric chemistry and governing processes removing N_r from the atmosphere is both complex and diverse, reflecting the many compounds involved and their different characteristics. To be able to relate these processes more effectively, the present paper provides an overview of the governing atmospheric processes of N_r in the atmosphere, with particular attention to the nature of the emissions and removal processes, including bi-directional exchange where this occurs. This review represents further develops and an extensive analysis of this topic (Hertel et al., 2011), as originally conceived in the European Nitrogen Assessment (2011). While the paper focuses on Europe, the general principles are applicable globally. For the purpose of this review, we have disregarded N₂O as it has no health effects at ambient concentration levels and an insignificant role in atmospheric deposition of N. It should be noted though, that it does play a role in the chemistry of the stratosphere and as a climate forcer (Pinder et al., 2012), as well as in leading to depletion of stratospheric ozone (Brink et al., 2011; Butterbach-Bah et al., 2011).

2 Emissions

N_r is mainly released to the atmosphere as NH₃, NO_x, and in poorly quantified amounts of organic N such as amines. By comparison, emission of N₂O accounts for a much smaller fraction, representing about 15% of the total European N_r emission to the atmosphere (Leip et al., 2011).

2.1 Ammonia emissions

This section is devoted to the ammonia emissions and contains sub-sections regarding: Animal houses and manure storages, Manure application and mineral fertilizer, Grazing animals, Non-agricultural sources, Spatial distribution, and Long term trends.

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

In western countries, agricultural activities contribute 85 %–98 % of the atmospheric NH₃ emissions (Anderson et al., 2003; Gyldenkærne et al., 2005; Sutton et al., 2000a). These emissions arise in some cases from the direct breakdown of fertilizer, but more importantly from the breakdown of organic matter (like urea) within animal wastes.

The emission process is a chemical and physical process which is highly temperature dependent and varies significantly over day and season (Gyldenkærne et al., 2005; Skjøth et al., 2004). The agricultural sources may be grouped as:

- point sources i.e. animal houses, manure storage and “slurry lakes” (Sommer et al., 2006);
- application of manure and mineral fertilizer to fields (Smith et al., 2009);
- grazing animals (Petersen et al., 1998);
- other sources including releases from plants (Larsson et al., 1998) such as legumes (Gyldenkærne et al., 2005) and vegetation during management and senescence (Sutton et al., 2000b, 2009a, b).

And the non-agricultural sources include:

- wild animals (Anderson et al., 2003; Riddick et al., 2012; Simpson et al., 1999; Sutton et al., 2000a; Theobald et al., 2006);
- catalyst processes, mainly related to road traffic (Kean et al., 2009; van Vuuren et al., 2011);
- manufacturing processes such as production of fertilizer, glass wool, catalysts and cement;
- humans, pets and sewage systems (Reche et al., 2012; Sutton et al., 2000a);
- other sources such as landfill and non-anthropogenic sources including natural fires from ecosystems (Andreae and Merlet, 2001; Yokelson et al., 2007);

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- emission from plants due to compensation points (Farquhar et al., 1980; Husted et al., 2000; Massad et al., 2010a; Mattsson et al., 2009; Sutton et al., 1995);
- emission from sea surfaces (Barrett, 1998; Sørensen et al., 2003).

The above split of the agricultural sources has been used in a parameterization currently considered as the most advanced description of temporal and spatial variations in NH₃ emissions (Pinder et al., 2007). This parameterization was originally developed for Denmark (Gyldenkerne et al., 2005; Skjøth et al., 2004), later extended to Northern Europe (Skjøth et al., 2011), and is currently being further developed in an open source version within the FP7 project ECLAIRE (<http://www.eclair-fp7.eu>). This parameterization and other more simple approaches have been successfully implemented in the Atmospheric Chemistry and Deposition (ACDEP) model (Skjøth et al., 2004), the DEHM model (Brandt et al., 2012, Skjøth et al., 2011), the EMEP MSC-W model (Fagerli et al., 2007; Simpson et al., 2012), the local scale models OML-DEP (Geels et al., 2012; Sommer et al., 2009), the SCAIL (Theobald et al., 2009) screening method developed for the UK, as well as most recently the local scale models AIRMOD and ADMS (Theobald, 2012). We here use the mechanistic description in this model as the basis for describing NH₃ emissions.

2.1.1 Animal houses and manure storages

Highly complex surface models have been developed for describing NH₃ emissions from agricultural buildings (Muck and Steenhuis, 1982; Olesen and Sommer, 1993; Oudendag and Luesink, 1998; Zhang et al., 1994) in order to help better manage emissions. However, in chemistry-transport modelling (CTM) a simplified parameterization based on wind speed and temperature has proven more practical (Skjøth et al., 2004):

$$E(t) = C \times T(t)^{0.89} \times V(t)^{0.26} \quad (1)$$

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where C is a constant related to the amount of N or TAN (Total Ammoniacal nitrogen = sum of NH_3 and NH_4^+) in the manure at a given time and location, $T(t)$ is the temperature as function of time, and V is the wind speed or the rate of ventilation. Formula (1) is used to distribute a known annual emission into shorter time steps. In Northern Europe, pig and poultry sheds are heated in winter and have reduced ventilation (Fig. 2). The annual emissions from manure storage vary with type of storage, and temporal variation follows ambient temperature (Gyldenkærne et al., 2005; Skjøth et al., 2004). In warm areas and during warm periods, emissions from buildings reflect outdoor temperature (Seedorf et al., 1998). Buildings with pigs and poultry have significant emissions also in cold periods, due to their warmer indoor temperatures, when outdoor storages and naturally ventilated cattle barns have low emission rates. Based on these principles and ambient temperature, it is possible to simulate temporal variations in emissions from manure storage, pig, poultry and cattle sheds (Fig. 3).

2.1.2 Manure application and mineral fertilizer

NH_3 emissions from field application of manure and mineral fertilizer take place at distinct times of year and with relatively short duration compared with animal related point sources. Soil type (Loubet et al., 2010) and the application method are crucial for establishing the magnitude of the emission. Broad-spread application over the whole of the ground surface leads to very high emissions, whereas direct soil injection or band-spreading methods strongly reduce emissions (Skjøth et al., 2008). National regulations can significantly affect seasonality when these affect application methods (Skjøth et al., 2008) although these emissions have substantial uncertainties (Sintermann et al., 2012). In many countries manure application is constrained by almost no regulation, whereas in most Northern European countries, such application is banned during winter to limit nutrient wash out, especially under the EU Nitrate Directive. To overcome shortage in storage capacity, farmers in Northern Europe often empty their tanks in autumn to allow them to handle emissions from animals kept indoors throughout the winter. This practice leads to late autumn emission peaks in NH_3 emission. Also for

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mineral fertilizer there are regional differences, as farmers in Southern Europe start fertilizer application earlier than Northern European farmers.

2.1.3 Grazing animals

The time that animals are in the field, together with the N content in the grass, govern the emission from grazing animals. Animals that are feeding on grass with a high N content excrete large amounts of N (as TAN) compared with animals on grassland with less nitrogen rich grass (Petersen et al., 1998). However, urine from grazing animals quickly enters the soil (Webb et al., 2005) which reduces the emission in comparison e.g. to surface applied slurry. The larger dry matter content of slurries limits infiltration into the soil, which is the main reason for the larger emissions compared with urine from grazing animals. In Southern Europe, animals in general are in the field most of the year. In Northern Europe, only sheep stay in the field most of the time. In most countries cattle are inside sheds approximately half of the year (Skjøth et al., 2011). In principle temperature should increase NH_3 emission related to grazing animals or outdoor yards used by livestock, however, clear temperature responses have not always been seen in practise (Misselbrook et al., 2001). However studies suggest that emissions from the urine fraction are larger during summer than during autumn (Ryden et al., 1987).

2.1.4 Non-agricultural sources

Non-agricultural sources of NH_3 are generally poorly described, which is largely a consequence of the limited number of data on many different processes. These sources include: human sweat, excreta from pets and wild animals, exhaust from gasoline vehicles with catalytic converters, stationary combustion sources and industry, as well as evaporation from waste deposits (Sutton et al., 2000a). Elevated NH_3 levels have been reported near roads and in urban areas as a consequence of NH_3 emissions from vehicles fitted with catalytic converters (Cape et al., 2004). Evaporation from NH_4^+ containing salts is a possible source of NH_3 in Southern and Central European cities

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where strongly elevated levels have been found: Barcelona (Reche et al., 2012), Rome (Perrino et al., 2002) and Rijeka (Croatia) (Alebic-Juretic, 2008). Perhaps the best described non-agricultural sources of NH_3 are from colonies of wild seabirds (Blackall et al., 2007; Riddick et al., 2012; Wilson et al., 2004) and seals (Theobald et al., 2006). Although these contribute to a small fraction of global emissions, they occur in otherwise clean remote environments, representing major point sources under purely climatic control for the given animal densities.

2.1.5 Spatial distribution

EMEP (the Co-operative Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe: <http://www.emep.int>) and CORINAIR (CORE Inventory AIR emissions) have compiled inventories at the European level on annual mean emissions with a spatial resolution of $50 \text{ km} \times 50 \text{ km}$ (Tørseth et al., 2012). EDGAR (Emissions Database for Global Atmospheric Research, <http://edgar.jrc.ec.europa.eu>) and GEIA (Global Emissions Inventory Activity, <http://geiacenter.org>) databases are available on $0.1 \times 0.1^\circ$ resolutions, and EUROTRAC (EUREKA project on the transport and chemical transformation of trace constituents in the troposphere over Europe) GENEMIS (GENeration and Evaluation of eMISSion data; <http://www.gsf.de/eurotrac>) compiled emission inventories for the year 1994 with a grid resolution of $16.67 \text{ km} \times 16.67 \text{ km}$ (Schwarz et al., 2000; Wickert et al., 2001). The GENEMIS data have in some cases been used to redistribute EMEP emission to higher resolution also for subsequent years assuming unchanged relative distributions (Hertel et al., 2002). The need for high resolution inventories has been recognized by the model groups within MACC and MACC-II research programmes (<http://www.gmes-atmosphere.eu/>). They have therefore adapted a high resolution inventory with 7 km resolution of annual emissions at SNAP level-1 (Pouliot et al., 2012). The TNO MACC-II inventory is a combination of the officially reported inventories, information on geographic location of point sources and a correction procedure for inconsistencies. Figure 4 shows the spatial distribution of the EMEP and EDGAR emission data sets. The EDGAR data set has

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a considerably higher spatial resolution compared to the EMEP data set. The EDGAR data set is constructed centrally using a uniform methodology and uniform data set, while the EMEP data set is a compilation of national inventories that originates from national experts. One result is that the two inventories do not agree with respect to amount of emissions within national boundaries. Such differences can be substantial (de Vries et al., 2011a; de Vries et al., 2011b). Here these differences are clearly seen in Fig. 4, with considerably higher emissions in central Europe in the EDGAR data set compared to the EMEP data set. Emissions are the most important input to all CTMs (Simpson et al., 2011), but for more than a decade been considered among the largest uncertainties in these models (Reis et al., 2009). The differences between the different emission data sets can therefore be expected to have a direct impact on model results for both process descriptions and impact assessments. Whereas emissions are usually reported as annual totals, there is considerable variation on hourly, daily and seasonal time-scales which is represented poorly (if at all) in the data used for CTMs (Menut and Bessagnet, 2010; Pouliot et al., 2012).

2.1.6 Long term trends

NH₃ emissions have been reduced in countries like Denmark, Germany and the Netherlands, whereas for France, Sweden and Norway only minor changes have occurred over the past 15 yr, and in countries like China and the US emissions have increased (Fig. 5). The uncertainties in these trends, including the small response of NH₃ concentrations to reduced animal numbers in central Europe (Horvath and Sutton, 1998) and to emission abatement policies in the Netherlands and Denmark (Bjerrregaard, 2011; Bleeker and Sutton, 2006; Erisman et al., 1998; Skjøth et al., 2008; Sutton et al., 2003) have been a major topic of debate. The recent evidence (Bleeker et al., 2009; Horvath et al., 2009) indicates that increasing NH₃ concentrations in Eastern Europe were due to a reduction in SO₂ emissions (resulting from longer atmospheric lifetime of NH₃) a factor which also influenced the NH₃ trends in the Netherlands,

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Denmark and elsewhere. The observed trends in NH₃ concentrations are therefore a result of changes in NH₃ emissions as well as changes in atmospheric chemistry.

2.2 Nitrogen oxide emissions

This section is devoted to the nitrogen oxides emissions and contains sub-sections regarding: spatial distribution, temporal variation, trends and projections.

NO_x is generated at high temperatures in combustion processes mainly from oxidation of free atmospheric nitrogen (N₂). Road transport and public power generation are by far the largest contributors to the emissions – for Europe these sources each contribute 20 % to 25 % of total NO_x emissions (Vestreng et al., 2008). Industrial production, other mobile sources and shipping each contribute 10 % to 15 %, whereas smaller contributions are related to aviation, and biomass and agricultural burning (2 % to 3 %). Emissions from forest and agricultural soils are discussed in Sect. 2.3. Emissions from waste incineration and lightning can probably be neglected (< 1 %).

2.2.1 Spatial distribution

Annual inventories of NO_x emissions are available from similar sources as for NH₃ (Sect. 2), which includes EMEP, EDGAR, EUROTRAC, GENEMIS and TNO-MACC. In the same way as for NH₃, the differences between the EDGAR and EMEP NO_x inventories can be seen in Fig. 6. The major road networks and urban areas are obvious hotspots and therefore easily visible. The same is true for shipping, where main transport routes are easily detectable and emissions may overall be higher than in urban areas. As a result of European regulation, power plants and industry release pollutants at a height that leaves local urban areas almost unaffected by these emissions, although they naturally contribute to pollutant levels further downwind.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2.2 Temporal variation

Road traffic follows very regular temporal patterns. In Northern Europe the typical urban road traffic pattern include two rush hour periods – in the morning and afternoon. Often the morning rush hour is more pronounced than the afternoon rush hour, and this is reflected in emissions and resulting concentrations of NO_x, as e.g. reported for Denmark (Hertel et al., 2008). In some parts of Southern Europe there may even be four rush hours as people are off work in the middle of the day. Furthermore, different patterns are observed for weekday vs. weekend, and the pattern also differs for passenger cars and commercial vehicles. Furthermore, highways are used for transport over longer distance and these have thus different diurnal traffic pattern compared with urban streets. Seasonal variations are less pronounced compared with diurnal variations, although increased power generation for heating, and increased direct fuel use lead to increased NO_x emissions in winter in Northern Europe, whereas increased demand for air conditioning in summer is expected to lead to increased NO_x emissions in summer in Southern Europe.

2.2.3 Trends and projections

European NO_x emissions have followed a steady downward trend over the time period 1990 to 2005. In total, a reduction of 34 % has been seen for the 27 Member States of the European Union (EU27) as a result of regulation of emissions – like the EC Large Combustion Plant Directive and the EURO regulations for road traffic vehicles (Fig. 5). Over the next decades further reductions will result from stricter regulation of road traffic despite increasing vehicle numbers and increasing distance travelled (Vestreng et al., 2008). Phasing out of nuclear energy may, however, lead to additional coal burning with the result of increasing NO_x emissions from electricity generation. International shipping has been steadily increasing, as have the associated emissions, but reductions will result from implementation of the HELCOM convention that includes NO_x Emission Control Areas (NECA) (http://www.helcom.fi/Recommendations/en_GB/)

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rec28E_13/) for the Baltic Sea and the North Sea. In urban streets, NO₂ has not followed the decreasing trend observed for NO_x, (Carslaw et al., 2007; Lambrecht, 2007). The explanation is that the NO₂ to NO_x ratio in road traffic emissions has changed substantially in recent years as a result of late diesel technology vehicles and catalytic converters in gasoline driven vehicles that reduce the overall NO_x emission but with a higher NO₂ fraction in the remaining emission (Carslaw and Beevers, 2005). EU limit values for NO₂ are therefore currently exceeded in many European cities (Hertel and Goodsite, 2009), and this will persist until stricter EURO 6 norms are implemented from 2015 (http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=_CELEX:32007R0715:EN:NOT).

2.3 Emissions of NO from ecosystems and soil

In a global perspective, NO_x emissions from soil contribute > 40 % (Davidson and Kingerlee, 1997) (IGAC Newsletter Dec 2000), and > 10 % for some European countries (Butterbach-Bahl et al., 2004; Skiba et al., 1997; Stohl et al., 1996). Estimates of emissions of NO_x from agricultural soil varies by more than a factor of two and are mainly related to grazing animals and fertilizer (de Vries et al., 2011c). Emissions that are related to application of fertilizer are estimated to account for 40 % of the soil emissions (IGAC Newsletter, 2000) and up to 65 % for the USA (Hall et al., 1996). Rural agricultural areas receiving N fertilizers in countries with long dry periods are likely the largest sources of soil NO where they can be of the same order of magnitude as N₂O fluxes (Loubet et al., 2011). The NitroEurope Integrated Project (Sutton et al., 2007; <http://www.nitroeuropa.eu>) and the NOFRETETE (Nitrogen oxides emissions from European Forest Ecosystems) project point to European forests being large sources of NO from soils (Pilegaard et al., 2006) presumably affected by enrichment with N from atmospheric deposition (Butterbach-Bahl et al., 2011a, b).

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3 Transformations

In the following a description of the most important chemical transformation processes for N_r is provided with highlights concerning where these processes play a significant role.

5 3.1 NH_x chemistry in the atmosphere

New aerosol particles are formed in gas phase reactions between NH_3 and gas phase acids. However, NH_3 may also deposit onto existing atmospheric particles. Whenever sulphuric acid (H_2SO_4) is present, NH_3 practically always reacts with this compound in an irreversible process forming aerosol phase ammonium bisulphate (NH_4HSO_4) and ammonium sulphate ($(NH_4)_2SO_4$). Naturally this reaction depends on the availability of H_2SO_4 which is currently decreasing in concentrations in Europe. Early experiments showed that 50 % of the available NH_3 was converted in about 35 min at concentration levels in those days ($20 \mu g m^{-3} SO_2$ and $2.7 \mu g m^{-3} NH_3$) (Mckay, 1971). H_2SO_4 is formed from gas phase oxidation of SO_2 by hydroxyl (OH) radical or from aerosol phase conversion by hydrogen peroxide (H_2O_2) and ozone (O_3). The latter process is pH dependent, and may be catalysed by NH_3 , since uptake of NH_3 increases the pH of aerosols (ApSimon et al., 1994; Junge and Ryan, 1958).

The rate of conversion of gas phase SO_2 and NH_3 into aerosol phase ammonium sulphate has been studied in detail in laboratory experiments (Baldwin and Golden, 1979; Gupta et al., 1995; Huntzicker et al., 1980; McMurry et al., 1983). At high RH, the limiting factor for the transformation is the molecular diffusion of NH_3 to the acid particles, whereas at low RH only between 10 % and 40 % of the collisions between NH_3 gas molecules and H_2SO_4 -containing particles lead to reaction (Huntzicker et al., 1980; McMurry et al., 1983). For small particles, their relatively large surface area makes the diffusion process more efficient. It has, however, been shown that organic material on the surface of the particles may limit the uptake of NH_3 (Daumer et al., 1992). Whereas the NH_3 reaction with H_2SO_4 may generally be considered irreversible,

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



this is not always the case, and not the case for the reactions with other acid gases. While reaction to form ammonium sulphates is effectively not reversible when these exist in particulate form, the NH_3 can be returned back to the gas phase as a result of changes in atmospheric composition e.g. during cloud processing (Bower et al., 1995, 1997; Wells et al., 1997).

The presence of HNO_3 and/or hydrochloric acid (HCl) together with NH_3 leads to equilibrium between these gases and their aerosol phase reaction products – ammonium nitrate (NH_4NO_3) and ammonium chloride (NH_4Cl). For this reaction an equilibrium product, $k_{\text{eq}} = [\text{NH}_3][\text{HNO}_3]$ of the gas phase concentrations of NH_3 and HNO_3 at saturation of the air is applied. Experimental studies show that this may be expressed as a function solely of temperature and humidity (Stelson et al., 1979; Stelson and Seinfeld, 1982). Besides the reactions with H_2SO_4 and HNO_3 , NH_3 may also react with HCl and form NH_4Cl (Pio and Harrison, 1987; Raes et al., 2000). Whereas HCl is emitted as a primary pollutant from coal burning and waste incineration, HNO_3 is the main secondary pollutant from oxidation of NO_x .

Ice core data have indicated that the HCl input from sea salt de-chlorination has been enhanced by a factor of 2–3 during the second half of the 20th century similarly to the increase of the atmospheric acidity in response to growing NO_x and SO_2 anthropogenic emissions (Legrand et al., 2002). New measurement data indicate that in NW Europe, HCl concentrations are similar (in terms of mixing ratio) to those of HNO_3 in summertime. However, NH_4Cl concentrations are generally much lower than NH_4NO_3 concentrations. HCl is emitted from anthropogenic sources including domestic coal burning in Europe which was historically a significant source of NH_x in the atmosphere (Sutton et al., 2008), but also released in displacement reaction in sea spray particles when these take up HNO_3 (Wall et al., 1988). This displacement is most likely the explanation for HCl concentrations of up to 250 pptv observed in the marine boundary layer (Harris et al., 1992). The displacement plays an important role in halogen and O_3 cycling (van Glasow and Crutzen, 2007). The resultant decline in HNO_3 due to the reactions with sea salt can promote the dissociation of NH_4NO_3 , and this reaction has

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

therefore been given considerable scientific attention as e.g. in the EU projects ANICE (de Leeuw et al., 2001) and MEAD (Spokes et al., 2006). Model estimates suggest that equilibrium between HNO_3 and sea salt can take hours to days to reach steady state (Erickson et al., 1999; Keene et al., 2002), similar to time scales for transport across the coastal zone. Thus reactions may not be complete within the coastal zone and HNO_3 concentrations relative to NO_3^- can be highly variable depending on the history of the air mass and the history of the mixing between sea salt aerosols and polluted continental air (Fischer et al., 2006).

NO_3^- aerosol is found in the coarse as well as the fine mode as a result of the displacement reaction whereas NH_4^+ is present in the fine mode and only in very small amounts in the coarse mode (Yeatman et al., 2001) which affects the subsequent efficiency of deposition.

3.2 NO_y chemistry in the atmosphere

The chemistry of NO_y and O_3 in the atmosphere has been discussed in numerous reviews (see e.g. Crutzen et al., 1999; Jenkin and Clemitshaw, 2000), so only an outline is given here. A sketch showing the main chemical paths of NO_y is given in Fig. 7, and a short description including main research findings in this area is given in the following. The fraction of directly emitted NO_2 from road traffic in western countries has increased in recent years. However, in the tropospheric boundary layer the distribution between NO and NO_2 is governed to a large degree by O_3 that reacts quickly with NO to form NO_2 . In sunlight NO_2 photo-dissociates (wavelengths 290 to 420 nm) to form NO and the very short-lived oxygen ($\text{O}(^3\text{P})$) radical. The latter will in most cases again form O_3 in reaction with free oxygen (O_2), involving a third body (either an N_2 or O_2 molecule) that absorbs excess vibrational energy and thereby stabilizes the formed O_3 molecule. These reactions all have time-scales of seconds to minutes. The rate of reaction between NO and O_3 is temperature dependent, but has a typical value about $4 \times 10^{-4} \text{ ppbv}^{-1} \text{ s}^{-1}$. Under typical atmospheric boundary layer conditions, this reaction

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

will either lead to the complete conversion of all the O_3 to NO_2 , or to the conversion of all NO to NO_2 (Clapp and Jenkin, 2001). In a highly polluted atmosphere (e.g. an urban area) or close to pollution sources, the former behaviour is usually observed because, although O_3 is widely distributed in the lower atmosphere, its concentration is not usually high compared with NO in the highly polluted atmosphere, and hence O_3 concentrations become rapidly depleted. During daylight, the main fate of NO_2 is to undergo photolysis, which is reforming O_3 and NO (Dickerson et al., 1982). This photolysis reaction has a typical rate coefficient under summer conditions in the mid-afternoon at mid-altitudes of about $7 \times 10^{-3} \text{ s}^{-1}$. The NO - NO_2 - O_3 chemistry in urban streets may be simplified (Palmgren et al., 1996), assuming that O_3 is reformed instantaneously, and thereby disregarding the formation of $\text{O}(^3\text{P})$. This is done in one of the most commonly applied street pollution models – OSPM (Berkowicz, 2000; Kakosimos et al., 2010).

The OH radical initiates the oxidation of a wide range of compounds in the atmospheric boundary layer. OH interacts with peroxy radicals that are responsible for the formation of excess concentrations of photo-oxidants like O_3 . In the background troposphere, carbon monoxide (CO) plays a role in this system. In the reaction with OH radical CO is oxidized to CO_2 , and at the same time a hydroperoxy radical (HO_2) is formed. Whenever NO is present, the most important atmospheric reaction of the hydroperoxy radical (HO_2) radical is the conversion of NO to NO_2 . The hydroperoxy radical is one of many peroxy radicals that take part in the conversion of NO to NO_2 . Organic peroxy radicals (RO_2) are likewise important and are mainly formed by the attack of the OH radical on the organic compounds ubiquitously present in the polluted atmosphere. These reactions follow a similar path as the CO oxidation, leading to formation of alkyl peroxy radicals (Jenkin and Clemitshaw, 2000) that subsequently convert NO to NO_2 in the same way as the HO_2 radical. In combustion processes at high temperatures e.g. inside the engine of a petrol or diesel-driven vehicle, NO is formed from ambient N_2 . However, in very NO rich air e.g. inside the exhaust pipe of vehicles and inside emitting chimneys, a 3rd order reaction can take place between two NO molecules

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and O_2 with a reaction rate coefficient of $2.3 \times 10^{-38} \text{ cm}^6 \text{ molecules}^{-2} \text{ s}^{-1}$ (Hampson and Gavin, 1978) to give NO_2 . The further transformation of NO_2 to HNO_3 takes place with a typical rate of about 5 % per hour in the troposphere. The hydroxyl radical (OH) is formed in daytime in the presence of sunlight (Jenkin and Clemitshaw, 2000). The photo-dissociation of O_3 leads to the formation of both $\text{O}(^3\text{P})$ and $\text{O}(^1\text{D})$ radicals; a fraction of the latter reacts with water vapour to form two OH radicals. This reaction is in competition with $\text{O}(^1\text{D})$'s reaction with 3rd body O_2 or N_2 molecules to form $\text{O}(^3\text{P})$, that in turn reforms O_3 . The OH radicals initiate most of the degradation of hydrocarbons in the atmosphere, a chain of reactions that e.g. lead to the formation of high O_3 concentrations during summer.

In the night-time the nitrate (NO_3) radical has a less important but still somewhat similar role for the degradation of hydrocarbons in the atmosphere as the OH radical in daytime (for an extensive review of the chemistry, see Wayne et al., 1991). Despite the considerably lower reactivity compared with OH , its higher peak concentrations in the night time troposphere allow the NO_3 radical to play a major role in the transformation of organic compounds. The NO_3 radical is formed during night-time in reaction with NO_2 . Dinitrogen pentoxide (N_2O_5) is a reservoir compound for the NO_3 radical at low temperatures, but it is broken down to its precursors NO_2 and NO_3 at higher temperatures in the dark. The typical night time NO_3 radical concentrations in the atmospheric boundary layer are of the order 10^7 to $10^8 \text{ molecules cm}^{-3}$ (which is the pptv range), although both measurements and modelling suggest that values away from the surface may reach 100s of ppt (e.g. Brown et al., 2006; Riemer et al., 2009). During the daytime both NO_3 and N_2O_5 photo-dissociate so fast that the concentrations of these compounds are insignificant. In the tropospheric boundary layer the photolysis of NO_3 radical (with a typical noon lifetime of about 5 s) follows two different wavelength dependent paths forming NO and O_2 (wavelength $> 700 \text{ nm}$) or NO_2 and $\text{O}(^3\text{P})$ (wavelength $> 580 \text{ nm}$). Close to pollution sources from combustion processes e.g. road traffic or power plants, the NO_3 radical is quickly removed by reaction with NO leading to formation of two NO_2 molecules. In urban areas, emissions of NO rapidly destroy

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

NO₃ close to ground-level. However, above the surface at night NO₂ and O₃ can exist together in the absence of NO (at it has all been converted to NO₂), leading to NO₃ formation, as observed on a tower in London (Benton et al., 2010). During night time the heterogeneous conversion of N₂O₅ to HNO₃ is an important process. The lifetime of N₂O₅ with respect to this removal is of the order of minutes in the tropospheric boundary layer. This production of HNO₃ may in winter be equally important as daytime conversion of NO₂ by OH radical. As already described, particulate nitrate (NO₃⁻) is formed when HNO₃ reacts with NH₃ and forms new aerosols, and when it sticks to existing particles in the atmosphere. In addition organic NO₃⁻ may be formed from gaseous NO₂ on the surfaces of aerosols in other heterogeneous reactions. The NO₃ radical attacks alkanes by hydrogen abstraction in a similar way as the reactions of the OH radical, and again leading to formation of a peroxy radical (RO₂) that may oxidise an NO molecule to NO₂. Also for alkenes, the attack of the NO₃ radical is similar to the reactions of the OH radical; the NO₃ radical adds to the double bond. This reaction is followed by rapid O₂ addition which leads to the production of a peroxy radical. Reaction of soil emissions of NO with atmospheric OH has been suggested to provide an in-canopy source of HNO₃ (Farmer et al., 2006). HNO₃ concentrations usually peak during the day, regulated by the emissions of NO_x, photochemical activity and the gas/aerosol equilibrium of NH₄NO₃ shifting towards the gas phase with increasing temperature and decreasing relative humidity (see e.g. Andersen and Hovmand, 1994).

A reaction similar to reaction between NO₂ and OH forming HNO₃, but less important, is the reaction between NO and OH radical that leads to formation of nitrous acid (HONO). During daytime HONO photo dissociates (λ < 400 nm) rapidly back to the reactants. Therefore, HONO formed in the late evening may serve as a night-time reservoir of OH and NO, which are subsequently liberated again the following morning in the sunlight. Studies in the highly polluted Po Valley in Northern Italy have shown an interrelation between simultaneous peaks in NO_x concentrations and aerosol surfaces and peak HONO concentrations during foggy periods (Notholt et al., 1992). This

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



was taken as evidence for heterogeneous conversion on aerosol surfaces. Probably this type of conversion plays an important role also in many urban areas over Europe, but so far only few studies have been carried out. Similar heterogeneous reactions can also occur on land surfaces (Harrison et al., 1996), and daytime formation of HONO can take place due to photo-sensitised reduction of NO₂ on humid acid in soil (Stemmler et al., 2006). A recent study furthermore suggest that nitrate in agricultural soil may lead to emission of HONO (Su et al., 2011).

3.3 Organic nitrogen compounds in the atmosphere

The presence of atmospheric organic N compounds has been evident for years (Cape et al., 2011; Cornell et al., 2003; Neff et al., 2002), but direct measurements of individual species (except PAN, discussed below) are rather sparse. The recent evidence for organic N comes from analysis of precipitation samples for total N and comparison with inorganic N content, to give “dissolved organic N” (DON) by difference. This approach has been prone to several analytical artefacts (Cape et al., 2001), but DON may in fact contribute up to half of total water-soluble N in precipitation.

The fraction of N in precipitation in organic form depends highly on location, and on whether air masses are of marine or terrestrial origin. DON has been large ignored in estimating environmental consequences of N deposition; such ecological impacts of DON depend on the bioavailability of the organic N, and there is reason to believe that many, if not all, components of DON are biologically available (Krab et al., 2008; Lipson and Nasholm, 2001; Paerl and Whittall, 1999; Qualls and Haines, 1992).

DON appears to have multiple sources, including agricultural activities and many reduced N compounds of biological origin, e.g. urea, amines, amino acids etc. (Cape et al., 2011). There appears to be a DON contribution from marine air masses (Cornell et al., 1995, 2001; Cornell, 2011), and DON proportions are consistently high in unpolluted air, especially in the tropics. For continental/terrestrial samples, annual average concentrations of DON in precipitation appear to correlate with total N and are better correlated with NH₄⁺ than with NO₃⁻ (Zhang et al., 2012). The overall correlation

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



suggests a significant anthropogenic source of organic N either directly or via atmospheric reactions, and the better correlation with NH_4^+ than with NO_3^- support the suggestion of the importance of agricultural sources of DON emission to the atmosphere (Cape et al., 2011).

5 The seasonal variations in DON concentrations are not always correlated with NH_4^+ concentrations, implying that different sources are involved (Cape et al., 2004). There is limited evidence available from sampling of air directly using a nebulising mist sampler that both gas-phase and particle phase components contribute to water-soluble organic N (WSO_N) in the atmosphere, which leads to the occurrence of DON in precipitation
10 (Benitez et al., 2010). Organic N has been measured in fog (Zhang and Anastasio, 2001) and cloud water (Hill et al., 2007), but there is some concern that most analyses for DON are made on bulk rainfall samples (i.e. collected using an open funnel) and that a significant fraction of the measured DON might have been dry-deposited on the funnel surface (Benitez et al., 2010). This presents problems of interpretation, but does
15 not remove the problem of identifying the source, composition and fate of organic N compounds.

Some measurements of individual components of reduced organic N in gas, particulate and aqueous phases have been reported and indicate potential sources and fates of these compounds, but in most cases rather small concentrations are measured and these cannot account for the rather high proportions of DON in precipitation. Recent
20 very sophisticated analyses of the composition of DON aimed at identifying functional groups are consistent with an important source involving reduced N, probably via reactions in the atmosphere (Altieri et al., 2012). Correlations between total N and DON in global databases and the remote atmosphere suggest DON represents on average
25 about 25 % of atmospheric N deposition, although this proportion is highly variable in space and time.

Organic nitrogen compounds are also formed as secondary products in the atmosphere from the reaction of various organic compounds with oxidized nitrogen species (e.g. Jenkin and Clemitshaw, 2000). For example, when aldehydes are photo

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

dissociated or react with OH, an alkyl radical is formed, which in turn may form peroxy alkyl nitrates that serve as important reservoirs of NO₂ (Fig. 8). The most abundant of these nitrates is the peroxyacetyl nitrate (PAN). PAN is thermally unstable and equilibrium between peroxy acetyl radical and NO₂ on one side and PAN on the other side is established in the boundary layer. High PAN and O₃ concentrations are often observed together during photo chemical smog episodes. Thermal degradation gives PAN a lifetime of ~ 1.7 h at 273 K and 50 h at 263 K. The PAN formation competes with NO degradation of peroxy acetyl radical. This reaction is usually dominating at ppbv levels of NO meaning that PAN and other peroxy alkyl nitrates are usually formed only in the background atmosphere, but substantial PAN concentrations may still be observed in urban areas at relatively low temperatures. The peroxy alkyl nitrates include compounds produced in a similar way as PAN, but generated from biogenic isoprene emissions that may be of importance in southern Europe, and have similar thermal degradation pathways as PAN. PAN is an important atmospheric N reservoir species, but it is probably not an important source of DON (Cornell et al., 2003). PAN deposition has been modelled in many studies (Wu et al., 2012), and the dry deposition is much faster than expected on the basis of its solubility, with dry deposition velocities in daytime up to 10 mm s⁻¹, so much is still unknown about the mechanisms of its atmospheric removal (Turnipseed et al., 2006).

4 Dry deposition and bi-directional fluxes

N_r is monitored in many regional networks across the world, such as the European EMEP programme (www.emep.int) that includes both long-term observations (Tørseth et al., 2012 and campaign studies Aas et al., 2012), the NitroEurope Integrated Project (NEU; www.nitroeuropa.eu; Sutton et al., 2007) the US National Atmospheric Deposition Network (NADP; http://nadp.sws.uiuc.edu/), the Acid Deposition Monitoring Network in East Asia (EANET; www.eanet.cc) and several others. However, these networks measure air concentrations rather than fluxes, and dry deposition is estimated

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



using inferential modelling approaches (Erisman et al., 1994; Erisman and Draaijers, 1995), such as using measured air concentrations and meteorology in combination with a dry deposition algorithm (e.g. Flechard et al., 2011; Skjøth et al., 2002) which are underpinned by often sparse databases of campaign based process studies with limited geographical coverage. This is partly due to the fact that instrumentation to measure fluxes of sticky compounds such as NH_3 , HNO_3 or HONO are expensive and labour intensive to operate. The measurement of each individual N_r compound is technically more challenging than that of CO_2 fluxes. Robust low cost flux measurement approaches are lacking, although recent development of a Conditional Time-Averaged Gradient (COTAG) method (Famulari et al., 2010) shows promise for wide-scale deployment over long periods for short vegetation.

A first European flux measurement network for reactive nitrogen compounds was established within the NitroEurope IP (2011) (Skiba et al., 2009; Sutton et al., 2007; Tang et al., 2009). This network applied at a three-tier approach, where selected N_r compounds were measured at a network of 13 supersites, using advanced micrometeorological flux measurement techniques. At a further 9 regional sites the novel COTAG systems combined with a low-cost time-integrated approach for measuring soil gas fluxes (System for Inert Gas flux Monitoring by Accumulation, SIGMA, Ambus et al., 2010) were deployed, while deposition was derived at a further 50+ “inferential sites” from concentration measurements, using inferential techniques (Flechard et al., 2011; Sutton et al., 2007; Tang et al., 2009).

A comprehensively detailed spatial coverage of N_r deposition can only be achieved through numerical modelling (e.g. Simpson et al., 2007, 2011). The gaseous N_r compounds usually making the biggest contribution to dry deposition are NH_3 , HNO_3 and NO_2 . Their relative contributions to N deposition depend on the pollution climate. In agricultural areas NH_3 may dominate the atmospheric N loading (e.g. Hertel et al., 2006), while in more industrial and urban areas HNO_3 and NO_2 may be more important. This pattern is seen even on the European scale; Simpson et al. (Simpson et al., 2007) showed that reduced nitrogen dominates dry-deposition of N to forests close to

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the main agricultural areas, whereas oxidised nitrogen dry-deposits over much larger transport distances. In addition, NH_3 deposition depends on the N status of the receiving surface, with fertilised vegetation and vegetation receiving high atmospheric N deposition inputs acting as a less efficient sink or even, in the case of fertilized vegetation, a net source of NH_3 (Loubet et al., 2009; Sutton et al., 1993). In wet regions, leaf cuticles frequently provide an efficient sink for water soluble gases (NH_3 and HNO_3), although the cuticular sinks for both gases can become saturated under very dry conditions, increasing the importance of stomata as an uptake pathway (Nemitz et al., 2004b).

4.1 The dry deposition process

Dry deposition is the direct uptake of gases or aerosols at terrestrial or marine surfaces. The dry deposition of gases and particles is a continuous process and governed by their air concentrations, turbulent transport processes in the boundary layer, the chemical and physical nature of the depositing species, and the biological and chemical capability of the surface to capture or absorb the species (e.g. Fowler et al., 2009).

In relation to deposition transport, the boundary layer may be considered to consist of two layers: the fully turbulent layer and the quasi-laminar layer. The quasi-laminar layer is introduced to quantify the way in which pollutant transfer differs from momentum transfer in the immediate vicinity of the surface (Hicks et al., 1987). In this layer, the transport is dominated by molecular diffusion. Once at the surface, the chemical, biological and physical nature of the surface determines the capture or absorption of the gases and particles. Deposition to water surfaces (oceans or fresh waters) may thus be very different from deposition to vegetated surfaces on land.

The deposition process may be considered as a series of resistances, by analogy with an electrical circuit (Monteith and Unsworth, 2008). The resistances refer to the transport processes through the various “layers” defined above: turbulent transfer (usually denoted R_a), quasi-laminar (R_b) and surface (R_c). For a complex surface with several potential absorption sinks (e.g. vegetation) the resistance R_c may be viewed as

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a network of parallel resistances, representing transfer to the external leaf surface, through stomata, to water on the surface, or through the canopy to the underlying soil surface. The total resistance (R_t) is the sum of all the series and parallel resistances ($R_a + R_b + R_c$), and is usually expressed in units of s m^{-1} . The inverse of the total resistance ($1/R_t$) is known as the deposition velocity (v_d), representing the deposition flux (F) normalized by the air concentration at height z (X_z), i.e. $v_d(z) = -F/X_z$ and has units of m s^{-1} (Fowler et al., 2001a; Hicks et al., 1987).

The turbulent transfer resistance (R_a) depends upon the height at which the deposition flux is measured, so the total resistance (R_t) and deposition velocity (v_d) also vary with height above the surface. The transfer flux (F) is defined as the product of the air concentration of a gas or particles at height z , multiplied by the deposition velocity at height z , and in the absence of competing chemical reactions (Sorensen et al., 2005), does not vary with height, provided that the air concentration is horizontally uniform.

The resistance formulation described above assumes that the concentration of the gas at the absorbing surface is zero (e.g. on the leaf cuticle or within plant stomata). Where this is not the case (see below) the effect can be described mathematically either as a decreased driving force for deposition (concentration difference between height z and the non-zero surface concentration) or as an increased surface resistance. The interpretation as a surface resistance in this context has the disadvantage that it is unable to simulate the bi-directional fluxes induced by non-zero surface concentrations.

The deposition velocity (v_d) is often reported as a constant even though it depends on a set of variables, e.g. wind speed, surface roughness and atmospheric stratification. Joffre (1988) has suggested a parameterization which depends on the meteorological conditions, roughness length and the molecular diffusion coefficient for the compound of interest. The various components of the total transport resistance can be estimated from meteorological data if several assumptions are made concerning spatial and temporal homogeneity (Erisman and Draaijers, 1995; Hicks et al., 1987; Nemitz and Sutton, 2004; Smith et al., 2000). The surface resistance (R_c) term depends on the physical and chemical nature of the absorbing surface, and parameterisations should

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

be adapted to the surface concerned. v_d is often expressed as annual or seasonal averages, for the purpose of calculating deposition fluxes as the product of air concentrations and deposition velocities.

When calculating fluxes, deposition velocities need to be used together with concentrations from the same height (z) – usually the height at which the concentrations are measured. Tall vegetation causes increased atmospheric turbulence, so R_a values are smaller, and deposition velocities are larger, than to short vegetation. Consequently, estimating deposition of different components requires knowledge about land-cover as well as the spatial pattern of air concentrations. The air-sea gas exchange of the very soluble gases HNO_3 and NH_3 is rate limited by the vertical transport in the boundary layer, because the uptake at the water surface is very fast relative to other commonly studied gases. Of the two very soluble N-gases, HNO_3 exchange rates tend to be generally larger than NH_3 due to the higher solubility (i.e. smaller R_c), although NH_3 deposition can be faster to a perfect sink due to its smaller molecular weight than HNO_3 and correspondingly smaller boundary layer resistance (R_b). The less soluble NO_2 and NO gases deposit much slower to the marine surface. The surface resistance is the most important resistance for slightly soluble gases and relates to the transfer velocity K_c , which is also used for air-sea exchange of other gases like CO_2 , DMS and CH_4 .

The surface resistance (R_c) is a key parameter for the deposition of a gas and several parameterization schemes have been derived from flux measurements. For sub-micron particles, the transport through the boundary layer is more or less the same as for gases. However, transport of particles through the quasi-laminar layer governing R_b can differ. For particles with an aerodynamic diameter $< 0.1 \mu\text{m}$, deposition is controlled by diffusion, whereas deposition of particles with an aerodynamic diameter $> 10 \mu\text{m}$ is more controlled by sedimentation (Seinfeld and Pandis, 2006). Deposition of particles with an aerodynamic diameter between 0.1 and $1 \mu\text{m}$ is determined by the rates of impaction and interception and depends heavily on the turbulence intensity. Transfer through the quasi-laminar layer close to the surface presents a considerable restriction on the deposition of 0.1 – $1.0 \mu\text{m}$ aerodynamic diameter particles. Uptake of particles

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

by surfaces is thus largely controlled by micro-structures and turbulence intensity. Although a range of theory-based models is available to describe aerosol deposition, they often predict features which conflict with measured deposition rates (Flechard et al., 2011; Petroff et al., 2008; Pryor et al., 2008a, b). For example, methods based on the well-known formulations of Slinn (Slinn, 1982) predict low deposition velocities to forest canopies. Alternative formulae of Zhang et al. (Zhang et al., 2001) predict higher deposition velocities, but no effect of canopy density. Several studies show that NH_4NO_3 has higher deposition velocities than sulphates, as a result of the partitioning of NH_4NO_3 to the more rapidly depositing HNO_3 and NH_3 gases (e.g. Fowler et al., 2009; Nemitz et al., 2004a; Wolff et al., 2010). Deposition of particles containing SO_4^{2-} , NO_3^- , Cl^- and NH_4^+ contributes to the potential acidification and eutrophication (N components) of ecosystems. Compared to gaseous deposition of acidifying compounds onto low vegetation, particle deposition fluxes are usually found to be small. However, unlike wet deposition it takes place all the time and furthermore it is believed that the fluxes of small particles are currently underestimated for very rough surfaces like forests. Erisman et al. (1996) found that deposition of aerosols to the Speulder forest contributed 20 and 40 % to the total dry deposition of S and N, respectively. Parameterisations of aerosol dry deposition velocities to forests differ greatly between models (Flechard et al., 2011).

4.2 Bi-directional fluxes of N-containing gases

Although reports of transient N_2O deposition fluxes are increasing in number, e.g. (Flechard et al., 2007), in terms of N inputs, the magnitude of N_2O uptake is small and negligible compared with the main contributors to atmospheric N_r deposition. For other N_r containing gases there are several parallel pathways of pollutant exchange with vegetation, including adsorption to leaf cuticles, exchange through the stomata with the sub-stomatal cavity and exchange with the soil. All these processes are potentially bi-directional, depending on the relative magnitude of the air concentration and the gaseous concentrations in chemical equilibrium with the leaf surface, the apoplastic

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fluid and the soil solution, respectively. The likelihood for uptake increases with the water solubility and Henry's Law coefficient of the gas, which vary several orders of magnitude (www.mpch-mainz.mpg.de/~sander/res/henry.html).

4.2.1 Nitric acid

5 Due to its high deposition rate, HNO_3 makes a significant contribution to Nr deposition in regions with high NO_x emissions. HNO_3 is highly water soluble and commonly assumed to deposit at the maximum rate permitted by turbulence, i.e. surface resistance is negligible. This is probably a reasonable approximation for most situations, but emission gradients or reduced uptake rates of HNO_3 have been observed, probably due to
10 non-zero HNO_3 surface concentrations in equilibrium with NH_4NO_3 deposited to leaf surfaces (Neftel et al., 1996; Nemitz et al., 2004; Zhang et al., 1995). For trace gases with negligible surface resistance, the deposition velocity is sensitive to the atmospheric resistances (R_a and R_b), which over rough surfaces are usually small ($5\text{--}10\text{ s m}^{-1}$). In such conditions, even a small surface resistance would strongly influence deposition
15 rates. Currently there are insufficient field data to show whether HNO_3 deposition is subject to a surface resistance, and this remains a research priority.

4.2.2 Ammonia

NH_3 is less water soluble than HNO_3 , and NH_3 previously absorbed to wet leaf surfaces may more readily be desorbed (re-emitted) as leaf water layers dry out again so
20 long as this has not yet be absorbed into the leaf tissues (Flechard et al., 1999; Sutton et al., 1998). Generally plants contain inorganic N in the form of NH_4^+ and NO_3^- . These nutrients are mainly present in the liquid part (apoplast) between the cells of the plant. NH_4^+ is an important by-product of plant biochemical pathways resulting in non-zero
25 NH_4^+ concentrations in the leaf apoplast, which results in non-zero gas-phase concentrations (stomatal compensation points, χ_s) in equilibrium with this $\text{NH}_{4\text{apo}}^+$ concentration

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

at the apoplastic pH (Farquhar et al., 1980; Massad et al., 2010b; Mattsson et al., 2009; Schjørring et al., 1998; Sutton et al., 1995).

Available evidence suggests that NH_4^+ increases with increasing N supply to the plant, either through fertilization or high atmospheric N inputs (Massad et al., 2010b; Sutton et al., 1995). The system is here an exchange governed by the pH depending relation between NH_3 and NH_4^+ . The compensation point χ_s is the product of a temperature function describing the Henry's Law equilibrium and the ratio of $\Gamma_s = [\text{NH}_4^+]_{\text{apo}} / [\text{H}^+]_{\text{apo}}$. Values of Γ_s range from < 100 for semi-natural vegetation in clean, remote environments to values around 1500 for semi-natural vegetation in environments with high N_r deposition, and to $> 10\,000$ after fertilization (Massad et al., 2010b). At 10°C , this equates to values of χ_s of < 0.15 , 2.3 and $> 15\,\mu\text{g NH}_3\text{ m}^{-3}$, respectively. Emission potentials of fertilized soils and plant litter can be several orders of magnitude larger (Nemitz et al., 2000; Sutton et al., 2009a, b), allowing NH_3 emitted at the ground to be recaptured by an overlaying plant canopy (Denmead et al., 1976). The decomposition of plant litter has been found to play an important role in agricultural canopies, such as oilseed rape and managed grassland (Denmead et al., 1976; Nemitz et al., 2000; Sutton et al., 2009b, c; Zhang et al., 2010) but is much less certain will often be less important for semi-natural vegetation, especially where litter has lower nitrogen content or is more acidic.

The large range of plant, litter and soil Γ values illustrates that the direction of NH_3 exchange is often difficult to estimate a priori. There are several reviews in the literature on bi-directional NH_3 exchange which have compiled data on compensation points (Massad et al., 2010b; Zhang et al., 2010) in order to provide the necessary input for application in atmospheric transport models. The compensation points increase with N input as it is the main driver of apoplast and bulk leaf NH_4^+ concentrations (Massad et al., 2010a; Sutton et al., 1993; Zhang et al., 2010), but the compensation point also varies between different plant species even when growing under the same conditions (Mattsson et al., 2009) and with growth stage and season (Milford et al., 2009; Riedo et al., 2002).

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The stomatal pathway for NH_3 exchange is only available when stomata are open during daytime, and thus deposition to (often wet) leaf surfaces is the dominant pathway during the night, unless soil surfaces provide a major source and are well exposed to the atmosphere. Maps of NH_3 net exchange are particularly uncertain, due to (i) uncertainties in the overall magnitude as well as spatial and temporal patterns of agricultural NH_3 emissions and (ii) the large variability of NH_3 deposition rates to different surfaces.

The development of resistance models to describe bi-directional NH_3 exchange for eventual incorporation in atmosphere transport models has progressed through several clear stages in recent decades. Initially, atmospheric transport models only allowed the application of uni-directional dry deposition, using values of v_d (Asman and Janssen, 1987). Introduction of the idea of a stomatal compensation point by Farquhar et al. (1980) then led to incorporation of simple compensation point values in atmospheric models (see Fischer, 1987). The next stage of development can with the introduction of the canopy compensation point approach, which offset bi-directional stomatal exchange with uptake onto leaf surfaces (Sutton et al., 1995; Sutton and Fowler, 1993), with this approach subsequently being adopted in both small- and large-scale atmospheric transport models by Dentener and Crutzen (1994); Sorteberg and Hov (1996) and Loubet et al. (2001). For example, Sorteberg and Hov (1996) found a reduction of 0–20 % in total sulphur deposition and a 0–25 % increase in NH_3 deposition when using a bidirectional scheme compared with a simple flux model allowing only NH_3 deposition. Lately Wichink Kruit et al. (2012) found that the inclusion of a compensation point model in the LOTUS-EUROS model redistributed N depositions on the regional scale and provided a considerable increase in NH_3 concentrations over sea areas, thus reflecting observed open sea concentrations (e.g. de Leeuw et al., 2003) to a much larger degree.

While the single layer canopy compensation point approach provided the foundation to simulate competing stomatal and cuticular processes within the plant canopy, it did not address the contribution of ammonia emissions from the ground surface or allow

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

dynamics of bidirectional cuticular exchange. The latter process was subsequently addressed, allowing simulation of NH_3 release from drying leaf surfaces, initially using a fixed cuticular pH (Neiryndk et al., 2005; Sutton et al., 1998), and subsequently extended to simulate full leaf surface chemistry, the model solving leaf surface pH according to comprehensive ion exchanges (Burkhardt et al., 2009; Flechard et al., 1999). Until now, this dynamic approach to treating leaf cuticle adsorption and desorption has been considered too complex for incorporation in atmospheric transport models.

By contrast to cuticular adsorption/desorption, the inclusion of a two-layer canopy compensation point approach has been considered more tractable for general model application. The basis of this approach was introduced by Nemitz et al. (2000, 2001), and has since formed the foundation for generalized empirical parameterizations estimated by Massad et al. (2010b) and Zhang et al. (2010) for use in atmospheric dispersion models e.g. Gore et al. (2009).

Experimental studies have shown that over the sea the atmospheric fluxes of NH_3 may also be upward or downward (Lee et al., 1998; Quinn et al., 1988; Sørensen et al., 2003) depending on the meteorological conditions and the relationship between the pH and NH_4^+ concentration in the upper surface waters on the one side, and the NH_3 concentrations in ambient air just above the water surface on the other side. The bi-directional NH_3 flux over sea is expressed as a simple compensation point exchange with the water surface: $F = V_e (C_{\text{eq}} - C_{\text{air}})$, where v_e is the exchange velocity between air and sea (that equals $1/(R_a + R_b)$), C_{eq} is the NH_3 concentration in the air at equilibrium with the NH_x in the water, and C_{air} is the actual ambient air concentration of NH_3 . The ambient air NH_3 concentration at equilibrium was expressed by Asman et al. (1994) as a function of NH_x concentration in sea water, activity coefficients for NH_3 and NH_4^+ in sea water, Henry's law coefficient for NH_3 and pH in sea water (Asman et al., 1994). The formulation has since been applied into the Lagrangian ACDEP model (Sørensen et al., 2003) using a distribution of NH_x concentrations in sea water from Barret (Barrett, 1998), where the results showed a redistribution of N deposition in the coastal region off the coast of the Netherlands. On a global basis Johnson et al. (2008)

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

emphasised the importance of temperature control on air-sea NH_3 exchange over the remote ocean and argued that NH_3 emissions were much more likely over warm waters than over cold (Johnson et al., 2008). Similarly, results with the ACDEP model suggested high sensitivity to both NH_4^+ concentration and pH of the sea water (Sørensen et al., 2003).

4.2.3 Nitrogen monoxide

NO is rather water-insoluble and there is no efficient mechanism for NO to react on the surface or inside leaves, so its deposition rate is rather slow. By contrast, soils are a source and sink for NO . Some of these soil emissions of NO are oxidized to NO_2 (and possibly HNO_3) within plant canopies, and taken up more efficiently than NO and thus the behaviour of NO still needs to be taken into account in surface – atmosphere exchange.

4.2.4 Nitrogen dioxide

Plant uptake of NO_2 is slower than that of the more water soluble gases (HNO_3 , NH_3), but it is a significant contributor to N deposition. The NO_2 deposition to vegetation is primarily regulated by stomata, and for most plants the internal resistance is negligible, and NO_2 deposition velocities may thus be computed from knowledge of stomatal resistance or conductance (Thoene et al., 1991). Studies indicate a small effective stomatal compensation point for NO_2 for some plant species, in the range of > 0 to 2 ppb; e.g. an American experimental study found a value of 1.5 ppb for the canopy compensation point for NO_2 over deciduous forest (Horii et al., 2004). However, the underlying process is not currently understood, and some laboratory work has failed to reproduce the field observations. Because of its low water solubility, deposition to (and reaction with) surface water, including sea water, is also slow (Cape et al., 1993).

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.2.5 Nitrous acid

The biosphere/atmosphere exchange of HONO is generally bi-directional, and day-time concentrations of HONO are low, as it is rapidly photolysed in sunlight. With solubility similar to NH₃, HONO is deposited to vegetation under most conditions. Observations of HONO emission have been attributed to production of HONO at surfaces, e.g. through the reaction of NO₂ with NO on wet surfaces (Harrison et al., 1994) or NO₂ reduction on humic acid (Stemmler et al., 2006). In connection with an experimental study, a parameterisation of bi-directional fluxes of both NH₃ and HONO was applied for estimating dry deposition of N to the Amazon basin from measured ambient air concentrations (Trebs et al., 2006).

4.2.6 Organic nitrogen compounds

Newly developed instruments have resulted in new measurements indicating deposition rates of PAN (and other PAN-like compounds) that are significantly larger than classical predictions (Turnipseed et al., 2006; Wolfe et al., 2009), especially to wet vegetation. Thus the lifetime of PAN with respect to deposition may be shorter than previously thought. In addition, PAN is water-insoluble and the comparably large deposition fluxes to wet surfaces indicate that the current mechanistic understanding of the deposition process is incomplete. There are parallels to the deposition of O₃, which also appears to exhibit larger deposition rates to wet surfaces than can be explained by its solubility (Fowler et al., 2001b). The importance of alkyl nitrates has recently been demonstrated for Blodgett Forest, Sierra Nevada, USA (Farmer et al., 2006), although it appears that the pollution climate of their site is unusual. Nevertheless, information is lacking to form a robust picture of the importance of these compounds across the full range of European conditions. Although amines have been measured as emitted from agricultural activities (Schade and Crutzen, 1995), there is currently no information on their dry deposition. Amines play a significant role in atmospheric new particle formation. Petäjä et al. (2011) showed that under atmospherically relevant conditions

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



amines are needed to explain cluster formation (Petäjä et al., 2011). Also several laboratory experiments have pointed out that amines have a marked enhancing impact on particle formation (Berndt et al., 2010; Murphy et al., 2007). Kurten et al. used quantum chemistry calculations and concluded that amines are far more efficient than ammonia in stabilizing small H_2SO_4 -clusters in the atmosphere (Kurten et al., 2008).

5 Wet deposition

Unlike dry deposition, the wet deposition processes are indirect in that rain, hail and snow are vectors for conveying gaseous and aerosol compounds to the surface. The simple precipitation collectors applied in monitoring networks contrast appreciably with the underlying physical and chemical pathways of solutes into the collected precipitation sample. There is also significant uncertainty in the relative magnitudes of dry deposition of trace chemical species as gases and aerosols onto the collecting equipment although this issue can be minimised in “wet only” collectors which are covered during dry periods.

5.1 Wet scavenging of aerosols

The bulk of the NH_4^+ and NO_3^- aerosol mass is present in the size range 0.1 to $1.0\ \mu\text{m}$ (diameter). These aerosols are removed through interception by falling rain or snow, a process known as washout or by incorporation of the aerosol into cloud droplets within clouds, a process known as rainout. Washout is responsible for 10 % to 20 % of the N_r in wet deposition. The aerosol scavenging within cloud occurs through a number of physical and chemical pathways while the gases are incorporated through solution and oxidation processes. The phoretic process includes diffusiophoresis, in which aerosol particles are transported in the direction of a mean flux of vapour molecules. In the case of a cloud droplet growing by vapour diffusion of water molecules towards the droplet surface, aerosols would move along the vapour flux towards the growing

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

droplet. Additional phoretic mechanisms are presented by electrical and thermal gradients (electrophoresis and thermophoresis respectively). The phoretic processes contribute relatively small amounts of the solute in cloud water (Goldsmith et al., 1963).

Aerosols may also be captured by cloud droplets following Brownian diffusion to the droplet surface and rates of Brownian diffusion vary strongly with particle size, being significant for particles smaller than 100 nm in diameter. However, diffusion rates are very small relative to molecular diffusion and diffusional mechanisms make only minor contributions to the wet removal pathway. The remaining minor process leading to capture of aerosols by cloud droplets is impaction and interception. As implied in the name these processes lead to the capture of aerosols by droplets when one is unable to follow the streamlines of airflow around the other and the aerosol and droplet collide. The bulk of the aerosol N in cloud water is incorporated through the activation of aerosols containing NO_3^- or NH_4^+ into cloud droplets. The N containing aerosols are effective cloud condensation nuclei and are readily incorporated into cloud droplets through the nucleation scavenging pathway. Thus the main route is nucleation scavenging for aerosol NO_3^- , and NH_4^+ (Pruppacher and Jaenicke, 1995). The pathway for below-cloud wet scavenging of the gaseous N_r depends on the solubility and reactivity of the specific gas. NH_3 and HNO_3 are highly soluble, and clouds and rain remove these gases effectively from the air. The contribution of NO and NO_2 to dissolved N in precipitation is very small as these gases are not very soluble. Wet deposition is monitored by simple methods (precipitation collectors) analysed for major anthropogenic ions SO_4^{2-} , NO_3^- , NH_4^+ , H^+ and marine ions Cl^- , Na^+ , Mg^{2+} . The networks of collectors for precipitation chemistry are much less dense than precipitation collectors for measurement of rainfall amounts, due to costs of chemical analysis. Furthermore, precipitation chemistry collectors are located at a height above ground to reduce contamination from ground based sources (Erisman et al., 2003), and the practice of locating collectors above the ground reduces the capture of small droplets due to aerodynamic screening by the collector (Dammgen et al., 2005).

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5.2 Distribution between dry and wet deposition

The relative contributions to deposition from dry and wet deposition change with distance from source as primary pollutant concentrations decline and oxidation from gas to particle removes gas phase species which dry deposit quickly. Thus areas more than a few hundred km from sources receive most of their N_r deposition in precipitation (Bartnicki et al., 2011; de Leeuw et al., 2003; Hertel et al., 2003), except possibly for forests whose aerodynamic roughness maximises dry-deposition (Simpson, 2006 1537/id). In regions in which the amounts of precipitation are large, wet deposition dominates the N_r loads, as in most of the uplands of Europe. However, it is not simply the precipitation amount that needs to be considered in assessing the relative contributions of wet and dry deposition. The processes leading to orographic enhancement of rainfall amount have a profound effect on the overall scavenging of pollutants from the atmosphere.

5.3 Orographic effects

The meteorological process which enhances precipitation in much of maritime northern Europe is the seeder – feeder mechanism, in which orographic cloud, formed over hills and mountains is washed out by precipitation falling from higher levels in the troposphere (Bergeron, 1965). Mountains are very effective in increasing rainfall and wet deposition by the seeder-feeder process in which low level hill cloud droplets are washed out by falling precipitation from higher levels. The hill cloud is more polluted than higher level cloud because boundary layer aerosols are effectively activated into cloud droplets as they are forced to rise and cool over the hills and mountains. The seeder – feeder effects on precipitation amount have been simulated in process-based models and are able to simulate observed spatial patterns in precipitation (Carruthers and Choularton, 1983). Models have also been used to simulate the wet deposition of pollutants over mountains (Dore et al., 1990) and compared with detailed campaign measurements in an upland area. Extending the modelling of orographic enhancement of wet deposition to the country scale has enabled detailed spatially resolved wet deposition maps to be

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

generated (Dore et al., 1990). As orographic enhancement of wet deposition has been shown to be a major contributor to the total deposition in upland Britain the explicit inclusion of the process in deposition maps has been regarded as a routine component of wet deposition mapping (<http://www.rotap.ceh.ac.uk>). The resulting wet, and total N deposition maps show a strong influence of altitude and require a grid resolution on the same scale as the complex topography to reproduce (< 10 km). Thus deposition modelling and mapping at a $50 \text{ km} \times 50 \text{ km}$ scale fails to capture the spatial structure in wet deposition (Fig. 8; Simpson et al., 2011). In principle the models are able to simulate the process, but the grid resolution of both the underpinning meteorological model and the model applied for deposition calculations needs to be able to capture the topographic scale of the variability and also take into account the spatial variation in the emissions (Dore et al., 2012), which for the UK area have been shown to be a spatial resolution of at least 1 km (Dore et al., 2012).

5.4 Cloud droplet deposition

Unlike aerosols in the size range $0.1\text{--}1.0 \mu\text{m}$, which are not deposited efficiently on vegetation, the hill cloud droplets are large enough (3 to $10 \mu\text{m}$ in diameter), to impact efficiently on vegetation (Fowler et al., 1990); this deposition pathway is termed cloud deposition or occult deposition. For the UK it provides a very small contribution to the total but it is important for hills which are frequently shrouded in cloud. As the concentrations of major ions in hill cloud are enhanced, this deposition pathway leads to the exposure of vegetation to very high concentrations (SO_4^{2-} , NH_4^+ , NO_3^- , $1000 \mu\text{eq l}^{-1}$ to $2000 \mu\text{eq l}^{-1}$) (Fowler et al., 1990). The orographic enhancement of wet deposition is not included in the assessments of wet deposition in all countries, and for regions with only small areas of upland, this will not lead to significant underestimates in wet deposition. However, for areas of Europe in which seeder – feeder scavenging of pollutant represents a substantial contribution to total deposition, it is important to simulate the

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

process in mapping regional wet deposition, to avoid underestimating wet deposition and exceedances of critical loads.

6 Conclusions

The emissions of N_r compounds are mainly related to releases of anthropogenic origin. These releases have typically strong diurnal and in some case also seasonal variations that relate to human activities (NO_x and NH_3) and climatic variables (mainly NH_3). N_r compounds are subject to rapid transformations in the atmosphere which profoundly affect their deposition rates. Thus NH_3 has a fast deposition rate but it also reacts with acids to form NH_4^+ salt aerosols which have low dry deposition velocities, but are liable to wet deposition. As a consequence, then NH_4^+ containing aerosols are readily capable of long-range transport. NO_x emissions are mainly in the form of NO , but the emissions from road traffic contain an important fraction of primary NO_2 which has increased in recent years. NO is rapidly converted to NO_2 , which in turn converts to HNO_3 either through daytime reaction with the OH radical, or by heterogeneous conversion processes at night. HNO_3 is subject to rapid deposition, but may also convert to NH_4NO_3 which is subject to long-range transport. NO_2 may also be converted by reaction on land and aerosol surfaces to $HONO$, which is capable of photolysis to form OH radicals. Ambient air concentrations of N_r compounds are generally fairly well (often within $\pm 20\%$ – 30%) reproduced by state-of-the-art models, while estimates of deposition are considerably more uncertain (often more than $\pm 50\%$). A significant part of the uncertainty in current CTMs is related to the fluxes with the surface, both emissions and deposition. Studies of fluxes of N_r are therefore in considerable need, especially for sensitive ecosystems. These flux studies need to include detailed field studies, parameterization, application and testing of models covering both the biosphere and the atmosphere such as CTMs. Furthermore, long-term trends show very different emission patterns for NH_3 and NO_x , where emissions in Europe and the US show marked declines for NO_x and moderate reductions or stagnation for NH_3 . In contrast, emissions

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from China, for instance, show a steep increase in the past decade and continue to grow rapidly (Reis et al., 2009). These variations will cause considerable changes in the atmosphere-biosphere system, where most N related feedback mechanisms remain to be studied (Arneth et al., 2010). Long-term studies of biosphere-atmosphere interactions in relation to N_f that take full cascade of effects into account is therefore highly needed.

Acknowledgements. This work was supported by the European Science Foundation programme Nitrogen in Europe (ESF-NinE), COST Action 729, the NitroEurope (The nitrogen cycle and its influence on the European greenhouse gas balance) Integrated Project (funded under the European Commission's 6th Framework Programme, contract 017841), and the ECLAIRE (Effects of Climate Change on Air Pollution and Response Strategies for European Ecosystems) (funded under the European Commission's 7th Framework Programme, contract 282910).

References

- Aas, W., Tsyro, S., Bieber, E., Bergström, R., Ceburnis, D., Ellermann, T., Fagerli, H., Frölich, M., Gehrig, R., Makkonen, U., Nemitz, E., Otjes, R., Perez, N., Perrino, C., Prévôt, A. S. H., Putaud, J.-P., Simpson, D., Spindler, G., Vana, M., and Yttri, K. E.: Lessons learnt from the first EMEP intensive measurement periods, *Atmos. Chem. Phys. Discuss.*, 12, 3731–3780, doi:10.5194/acpd-12-3731-2012, 2012.
- Alebic-Juretic, A.: Airborne ammonia and ammonium within the Northern Adriatic area, Croatia, *Environ. Pollut.*, 154, 439–447, 2008.
- Altieri, K. E., Hastings, M. G., Peters, A. J., and Sigman, D. M.: Molecular characterization of water soluble organic nitrogen in marine rainwater by ultra-high resolution electrospray ionization mass spectrometry, *Atmos. Chem. Phys.*, 12, 3557–3571, doi:10.5194/acp-12-3557-2012, 2012.
- Ambus, P., Skiba, U., Drewer, J., Jones, S. K., Carter, M. S., Albert, K. R., and Sutton, M. A.: Development of an accumulation-based system for cost-effective chamber measurements of inert trace gas fluxes, *Euro. J. Soil Sci.*, 61, 785–792, 2010.

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Andersen, H. V. and Hovmand, M. F.: Measurements of Ammonia and Ammonium by Denuder and Filter Pack, *Atmos. Environ.*, 28, 3495–3512, 1994.
- Anderson, N., Strader, R., and Davidson, C.: Airborne reduced nitrogen: ammonia emissions from agriculture and other sources, *Environ. Int.*, 29, 277–286, 2003.
- 5 Andreea, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cy.*, 15, 955–966, 2001.
- Aneja, V. P., Schlesinger, W. H., and Erisman, J. W.: Effects of Agriculture upon the Air Quality and Climate: Research, Policy, and Regulations, *Environ. Sci. Technol.*, 43, 4234–4240, 2009.
- 10 ApSimon, H., Barker, B. M., and Kayin, S.: Modeling Studies of the Atmospheric Release and Transport of Ammonia in Anticyclonic Episodes, *Atmos. Environ.*, 28, 665–678, 1994.
- Arneth, A., Harrison, S. P., Zaehle, S., Tsigaridis, K., Menon, S., Bartlein, P. J., Feichter, J., Korhola, A., Kulmala, M., O'Donnell, D., Schurgers, G., Sorvari, S., and Vesala, T.: Terrestrial biogeochemical feedbacks in the climate system, *Nat. Geosci.*, 3, 525–532, 2010.
- 15 Asman, W. A. H. and Janssen, A. J.: A Long-Range Transport Model for Ammonia and Ammonium for Europe, *Atmos. Environ.*, 21, 2099–2119, 1987.
- Asman, W. A. H., Harrison, R. M., and Ottley, C. J.: Estimation of the Net Air-Sea Flux of Ammonia Over the Southern Bight of the North-Sea, *Atmos. Environ.*, 28, 3647–3654, 1994.
- Baldwin, A. C. and Golden, D. M.: Heterogeneous Atmospheric Reactions – Sulfuric-Acid Aerosols As Tropospheric Sinks, *Science*, 206, 562–563, 1979.
- 20 Barrett, K.: Oceanic ammonia emissions in Europe and their transboundary fluxes, *Atmos. Environ.*, 32, 381–391, 1998.
- Bartnicki, J., Semeena, V. S., and Fagerli, H.: Atmospheric deposition of nitrogen to the Baltic Sea in the period 1995–2006, *Atmos. Chem. Phys.*, 11, 10057–10069, doi:10.5194/acp-11-10057-2011, 2011.
- 25 Benitez, J. M. G., Cape, J. N., and Heal, M. R.: Gaseous and particulate water-soluble organic and inorganic nitrogen in rural air in southern Scotland, *Atmos. Environ.*, 44, 1506–1514, 2010.
- Benton, A. K., Langridge, J. M., Ball, S. M., Bloss, W. J., Dall'Osto, M., Nemitz, E., Harrison, R. M., and Jones, R. L.: Night-time chemistry above London: measurements of NO_3 and N_2O_5 from the BT Tower, *Atmos. Chem. Phys.*, 10, 9781–9795, doi:10.5194/acp-10-9781-2010, 2010.
- 30

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bergeron, X.: On the low-level redistribution of atmospheric water caused by orography, in: the Proceedings of the International Conference on Cloud Physics, 24-05-1965, Tokyo and Sapporo, 96–100, 1965.

Berkowicz, R.: OSPM – A parameterised street pollution model, Environ. Monitor. Assess., 65, 323–331, 2000.

Berndt, T., Stratmann, F., Sipilä, M., Vanhanen, J., Petäjä, T., Mikkilä, J., Grüner, A., Spindler, G., Lee Mauldin III, R., Curtius, J., Kulmala, M., and Heintzenberg, J.: Laboratory study on new particle formation from the reaction OH + SO₂: influence of experimental conditions, H₂O vapour, NH₃ and the amine tert-butylamine on the overall process, Atmos. Chem. Phys., 10, 7101–7116, doi:10.5194/acp-10-7101-2010, 2010.

Bjerregaard, H.: 3.3 Impact assessment and regulation of N-emissions from livestock farms in Denmark in: Nitrogen Deposition and Natura 2000 – Science and practice in determining environmental impacts, edited by: Hicks, W. K., Whitfield, C. P., Bealey, W. J., and Sutton, M. A., COST, York, 38–45, 2011.

Blackall, T. D., Wilson, L. J., Theobald, M. R., Milford, C., Nemitz, E., Bull, J., Bacon, P. J., Hamer, K. C., Wanless, S., and Sutton, M. A., Ammonia emissions from seabird colonies, Geophys. Res. Lett., 34, L10801, doi:10.1029/2006GL028928, 2007.

Bleeker, A. and Sutton, M. A.: Linking ammonia emission reduction and air concentrations and depositions of reduced nitrogen in Europe, Abstr. Papers of the American Chemical Society, 232, 401, 2006.

Bleeker, A., Sutton, M. A., Achermann, B., Alebic-Juretic, A., Aneja, V., Ellermann, T., Erisman, J. W., Fowler, D., Fagerli, H., Gauger, T., Harlen, K. S., Hole, L. R., Horvath, L., Mitosinkova, M., Smith, R. I., Tang, Y. S., and Van Pul, A.: Linking ammonia emission trends to measured concentrations and deposition of reduced nitrogen at different scales in: Atmospheric Ammonia: Detecting emission changes and environmental impacts, edited by: Sutton, M. A., Reis, S., and Baker, S. M. H., Springer Press, 123–180, 2009.

Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., Van der Hoek, K. W., and Olivier, J. G. J.: A global high-resolution emission inventory for ammonia, Global Biogeochem. Cy., 11, 561–587, 1997.

Bower, K. N., Wells, M., Choularton, T. W., and Sutton, M. A.: A Model of Ammonia/Ammonium Conversion and Deposition in A Hill Cap Cloud, Q. J. Roy. Meteorol. Soc., 121, 569–591, 1995.

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Bower, K. N., Choularton, T. W., Gallagher, M. W., Colville, R. N., Wells, M., Beswick, K. M., Wiedensohler, A., Hansson, H. C., Svenningsson, B., Swietlicki, E., Wendisch, M., Berner, A., Kruisz, C., Laj, P., Facchini, M. C., Fuzzi, S., Bizjak, M., Dollard, G., Jones, B., Acker, K., Wieprecht, W., Preiss, M., Sutton, M. A., Hargreaves, K. J., Storetonwest, R. L., Cape, J. N., and Arends, B. G.: Observations and modelling of the processing of aerosol by a hill cap cloud, *Atmos. Environ.*, 31, 2527–2543, 1997.
- Brink, C., van Grinsven, H., Jakobsen, B. H., Rabl, A., Gren, I.-M., Holland, M., Klimont, Z., Hicks, K., Brouwer, R., Dickens, R., Willems, J., Termansen, M., Velthof, G., Alkemade, R., van Oorschot, M., and Webb, J.: Costs and benefits of nitrogen in the environment in: *The European Nitrogen Assessment – Sources, Effects and Policy Perspectives*, edited by: Sutton, M. A., Howard, C. M., Erisman, J. W., Billen, G., Bleeker, A., Grennfelt, P., van Grinsven, H., and Grizzetti, B., Cambridge University Press, Cambridge, 513–540, 2011.
- Brown, S. S., Ryerson, T. B., Wollny, A. G., Brock, C. A., Peltier, R., Sullivan, A. P., Weber, R. J., Dube, W. P., Trainer, M., Meagher, J. F., Fehsenfeld, F. C., and Ravishankara, A. R.: Variability in nocturnal nitrogen oxide processing and its role in regional air quality, *Science*, 311, 67–70, 2006.
- Burkhardt, J., Flechard, C. R., Gresens, F., Mattsson, M., Jongejan, P. A. C., Erisman, J. W., Weidinger, T., Meszaros, R., Nemitz, E., and Sutton, M. A.: Modelling the dynamic chemical interactions of atmospheric ammonia with leaf surface wetness in a managed grassland canopy, *Biogeosciences*, 6, 67–84, doi:10.5194/bg-6-67-2009, 2009.
- Butterbach-Bahl, K., Kesik, M., Miehle, P., Papen, H., and Li, C.: Quantifying the regional source strength of N-trace gases across agricultural and forest ecosystems with process based models, *Plant Soil*, 260, 311–329, 2004.
- Butterbach-Bahl, K., Nemitz, E., Zaehle, S., Billen, G., Boeckx, P., Erisman, J. W., Garnier, J., Upstill-Goddard, R. C., Kreuzer, M., Oenema, O., Reis, S., de Vries, W., Winiwarter, W., and Sutton, M. A.: Nitrogen as a threat to European air quality in: *The European Nitrogen Assessment – Sources, Effects and Policy Perspectives*, edited by: Sutton, M. A., Howard, C. M., Erisman, J. W., Billen, G., Bleeker, A., Grennfelt, P. I., van Grinsven, H., and Grizzetti, B., Cambridge University Press, Cambridge, 434–462, 2011a.
- Butterbach-Bahl, K., Gundersen, P., Ambus, P., Augustin, J., Beier, C., Boeckx, P., Dannemann, M., Gimeno, B., Kiese, R., Kitzler, B., Ibrom, A., Rees, R. M., Smith, K. A., Stevens, C. J., Vesala, T., and Zechmeister-Boltenstern, S.: Nitrogen processes in terrestrial ecosystems in: *The European Nitrogen Assessment – Sources, Effects and Policy Perspectives*, edited

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- by: Sutton, M. A., Howard, C. M., Erisman, J. W., Billen, G., Bleeker, A., Grennfelt, P., van Grinsven, H., and Grizzetti, B., Cambridge University Press, Cambridge, 99–125, 2011b.
- Cape, J. N., Storetonwest, R. L., Devine, S. F., Beatty, R. N., and Murdoch, A.: The Reaction of Nitrogen-Dioxide at Low Concentrations with Natural-Waters, *Atmos. Environ.*, 27, 2613–2621, 1993.
- Cape, J. N., Kirika, A., Rowland, A. P., Wilson, D., Jickells, T., and Cornell, S.: Organic nitrogen in precipitation: real problem or sampling artifact?, *The Scientific World*, 1, 230–237, 2001.
- Cape, J. N., Tang, Y. S., van Dijk, N., Love, L., Sutton, M. A., and Palmer, S. C. F.: Concentrations of ammonia and nitrogen dioxide at roadside verges, and their contribution to nitrogen deposition, *Environ. Pollut.*, 132, 469–478, 2004.
- Cape, J. N., Cornell, S. E., Jickells, T. D., and Nemitz, E.: Organic nitrogen in the atmosphere – Where does it come from?, *A review of sources and methods*, *Atmos. Res.*, 102, 30–48, 2011.
- Carruthers, D. J. and Choularton, T. W.: A Model of the Feeder Seeder Mechanism of Orographic Rain Including Stratification and Wind-Drift Effects, *Q. J. Roy. Meteorol. Soc.*, 109, 575–588, 1983.
- Carlsaw, D. C. and Beevers, S. D.: Development of an urban inventory for road transport emissions of NO₂ and comparison with estimates derived from ambient measurements, *Atmos. Environ.*, 39, 2049–2059, 2005.
- Carlsaw, D. C., Beevers, S. D., and Bell, M. C.: Risks of exceeding the hourly EU limit value for nitrogen dioxide resulting from increased road transport emissions of primary nitrogen dioxide, *Atmos. Environ.*, 41, 2073–2082, 2007.
- Clapp, L. J. and Jenkin, M. E.: Analysis of the relationship between ambient levels Of O₃, NO₂ and NO as a function of NO_x in the UK, *Atmos. Environ.*, 35, 6391–6405, 2001.
- Cornell, S., Rendell, A., and Jickells, T.: Atmospheric Inputs of Dissolved Organic Nitrogen to the Oceans, *Nature*, 376, 243–246, 1995.
- Cornell, S., Mace, K., Coeppicus, S., Duce, R., Huebert, B., Jickells, T., and Zhuang, L. Z.: Organic nitrogen in Hawaiian rain and aerosol, *J. Geophys. Res.-Atmos.*, 106, 7973–7983, 2001.
- Cornell, S. E., Jickells, T. D., Cape, J. N., Rowland, A. P., and Duce, R. A.: Organic nitrogen deposition on land and coastal environments: a review of methods and data, *Atmos. Environ.*, 37, 2173–2191, 2003.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Cornell, S. E.: Atmospheric nitrogen deposition: Revisiting the question of the importance of the organic component, *Environ. Pollut.*, 159, 2214–2222, 2011.
- Crutzen, P. J., Lawrence, M. G., and Poschl, U.: On the background photochemistry of tropospheric ozone, *Tellus Ser. A*, 51, 123–146, 1999.
- 5 Dammgen, U., Erisman, J. W., Cape, J. N., Grunhage, L., and Fowler, D.: Practical considerations for addressing uncertainties in monitoring bulk deposition, *Environ. Pollut.*, 134, 535–548, 2005.
- Daumer, B., Niessner, R., and Klockow, D.: Laboratory Studies of the Influence of Thin Organic Films on the Neutralization Reaction of H₂SO₄ Aerosol with Ammonia, *J. Aerosol Sci.*, 23, 315–325, 1992.
- 10 Davidson, E. A. and Kinglerlee, W.: A global inventory of nitric oxide emissions from soils, *Nutrient Cyc. Agroecosyst.*, 48, 37–50, 1997.
- Day, D. A., Wooldridge, P. J., and Cohen, R. C.: Observations of the effects of temperature on atmospheric HNO₃, ΣANs, ΣPNs, and NO_x: evidence for a temperature-dependent HO_x source, *Atmos. Chem. Phys.*, 8, 1867–1879, doi:10.5194/acp-8-1867-2008, 2008.
- 15 de Leeuw, G., Cohen, L., Frohn, L. M., Geernaert, G., Hertel, O., Jensen, B., Jickells, T., Klein, L., Kunz, G. J., Lund, S., Moerman, M., Muller, F., Pedersen, B., von Salzen, K., Schlunzen, K. H., Schulz, M., Skjøth, C. A., Sørensen, L. L., Spokes, L., Tamm, S., and Vignati, E.: Atmospheric input of nitrogen into the North Sea: ANICE project overview, *Cont. Shelf Res.*, 21, 2073–2094, 2001.
- 20 de Leeuw, G., Skjøth, C. A., Hertel, O., Jickells, T., Spokes, L., Vignati, E., Frohn, L., Frydendall, J., Schulz, M., Tamm, S., Sørensen, L. L., and Kunz, G. J.: Deposition of nitrogen into the North Sea, *Atmos. Environ.*, 37, S145–S165, 2003.
- de Vries, F. T., van Groenigen, J. W., Hoffland, E., and Bloem, J.: Nitrogen losses from two grassland soils with different fungal biomass, *Soil Biol. Biochem.*, 43, 997–1005, 2011a.
- 25 de Vries, W., Kros, J., Reinds, G. J., and Butterbach-Bahl, K.: Quantifying impacts of nitrogen use in European agriculture on global warming potential, *Curr. Opinion Environ. Sustain.*, 3, 291–302, 2011b.
- de Vries, W., Leip, A., Reinds, G. J., Kros, J., Lesschen, J. P., and Bouwman, A. F.: Comparison of land nitrogen budgets for European agriculture by various modeling approaches, *Environ. Pollut.*, 159, 3254–3268, 2011c.
- 30 Denmead, O. T., Freney, J. R., and Simpson, J. R.: Closed Ammonia Cycle Within A Plant Canopy, *Soil Biol. Biochem.*, 8, 161–164, 1976.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Dentener, F. J. and Crutzen, P. J.: A 3-Dimensional Model of the Global Ammonia Cycle, *J. Atmos. Chem.*, 19, 331–369, 1994.
- Derwent, R. and Hertel, O.: Transformation of Air Pollutants in: Urban Air Pollution – European Aspects, edited by: Fenger, J., Hertel, O., and Palmgren, F., Kluwer Academic Publishers, Dordrecht/Boston/London, 137–160, 1998.
- Dickerson, R. R., Stedman, D. H., and Delany, A. C.: Direct measurements of ozon and nitrogen dioxide photolysis rates in the troposphere, *J. Geophys. Res.-Atmos.*, 78, 4933–4946, 1982.
- Dore, A. J., Choularton, T. W., Fowler, D., and Stortonwest, R.: Field-Measurements of Wet Deposition in An Extended Region of Complex Topography, *Q. J. Roy. Meteorol. Soc.*, 116, 1193–1212, 1990.
- Dore, A. J., Kryza, M., Hall, J. R., Hallsworth, S., Keller, V. J. D., Vieno, M., and Sutton, M. A.: The influence of model grid resolution on estimation of national scale nitrogen deposition and exceedance of critical loads, *Biogeosciences*, 9, 1597–1609, doi:10.5194/bg-9-1597-2012, 2012.
- Erickson, D. J., Seuzaret, C., Keene, W. C., and Gong, S. L.: A general circulation model based calculation of HCl and ClNO₂ production from sea salt dechlorination: Reactive Chlorine Emissions Inventory, *J. Geophys. Res.-Atmos.*, 104, 8347–8372, 1999.
- Erisman, J. W. and Draaijers, G.: Atmospheric deposition in Relation to Acidification and Eutrophication, Elsevier, Amsterdam, 405 pp., 1995.
- Erisman, J. W., Beier, C., Draaijers, G., and Lindberg, S.: Review of Deposition Monitoring Methods, *Tellus Ser. B*, 46, 79–93, 1994.
- Erisman, J. W., Bleeker, A., and Van Jaarsveld, J. A.: Evaluation of ammonia emission abatement on the basis of measurements and model calculations, *Environ. Pollut.*, 102, 269–274, 1998.
- Erisman, J. W., Mols, H., Fonteijn, P., Geusebroek, M., Draaijers, G., Bleeker, A., and van der Veen, D.: Field intercomparison of precipitation measurements performed within the framework of the Pan European Intensive Monitoring Program of EU/ICP forest, *Environ. Pollut.*, 125, 139–155, 2003.
- Erisman, J. W., Galloway, J., Seitzinger, S., Bleeker, A., and Butterbach-Bahl, K.: Reactive nitrogen in the environment and its effect on climate change, *Curr. Opinion Environ. Sustain.*, 3, 281–290, 2011.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Fagerli, H., Hertel, O., Skjøth, C. A., Gyldenkerne, S., and Hutchings, N.: Implementation of a dynamical ammonia emission parameterization in the regional Unified EMEP chemical transport model Costa do Sauípe, Bahia, in Northeast Brazil, 1–5 October 2007.
- Famulari, D., Fowler, D., Nemitz, E., Hargreaves, K. J., Storeton-West, R. L., Rutherford, G., Tang, Y. S., Sutton, M. A., and Weston, K. J.: Development of a low-cost system for measuring conditional time-averaged gradients of SO₂ and NH₃, *Environ. Monitor. Assess.*, 161, 11–27, 2010.
- Farmer, D. K., Wooldridge, P. J., and Cohen, R. C.: Application of thermal-dissociation laser induced fluorescence (TD-LIF) to measurement of HNO₃, Σ alkyl nitrates, Σ peroxy nitrates, and NO₂ fluxes using eddy covariance, *Atmos. Chem. Phys.*, 6, 3471–3486, doi:10.5194/acp-6-3471-2006, 2006.
- Farquhar, G. D., Firth, P. M., Wetselaar, R., and Weir, B.: On the Gaseous Exchange of Ammonia Between Leaves and the Environment – Determination of the Ammonia Compensation Point, *Plant Physiol.*, 66, 710–714, 1980.
- Fischer, B. E. A.: Assessing recent ammonia inventories using a stationary long-range transport model, edited by: Asman, W. A. H. and Diederer, H. S. M. A., Bilthoven, The Netherlands, 1987.
- Fischer, E., Pszenny, A., Keene, W., Maben, J., Smith, A., Stohl, A., and Talbot, R.: Nitric acid phase partitioning and cycling in the New England coastal atmosphere, *J. Geophys. Res.-Atmos.*, 111, D23S09, doi:10.1029/2006JD007328, 2006.
- Flechard, C. R., Fowler, D., Sutton, M. A., and Cape, J. N.: A dynamic chemical model of bi-directional ammonia exchange between semi-natural vegetation and the atmosphere, *Q. J. Roy. Meteorol. Soc.*, 125, 2611–2641, 1999.
- Flechard, C. R., Ambus, P., Skiba, U., Rees, R. M., Hensen, A., van Amstel, A., Pol-van Dassel, A. V., Soussana, J. F., Jones, M., Clifton-Brown, J., Raschi, A., Horvath, L., Neftel, A., Jocher, M., Ammann, C., Leifeld, J., Fuhrer, J., Calanca, P., Thalman, E., Pilegaard, K., Di Marco, C., Campbell, C., Nemitz, E., Hargreaves, K. J., Levy, P. E., Ball, B. C., Jones, S. K., van de Bulk, W. C. M., Groot, T., Blom, M., Domingues, R., Kasper, G., Allard, V., Ceschia, E., Cellier, P., Laville, P., Henault, C., Bizouard, F., Abdalla, M., Williams, M., Baronti, S., Berretti, F., and Grosz, B.: Effects of climate and management intensity on nitrous oxide emissions in grassland systems across Europe, *Agr. Ecosyst. Environ.*, 121, 135–152, 2007.
- Flechard, C. R., Nemitz, E., Smith, R. I., Fowler, D., Vermeulen, A. T., Bleeker, A., Erisman, J. W., Simpson, D., Zhang, L., Tang, Y. S., and Sutton, M. A.: Dry deposition of reactive

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Geels, C., Andersen, H. V., Ambelas Skjøth, C., Christensen, J. H., Ellermann, T., Løfstrøm, P., Gyldenkerne, S., Brandt, J., Hansen, K. M., Frohn, L. M., and Hertel, O.: Improved modelling of atmospheric ammonia over Denmark using the coupled modelling system DAMOS, *Biogeosciences*, 9, 2625–2647, doi:10.5194/bg-9-2625-2012, 2012.



Goldsmith, P., Delafield, H. J., and Cox, L. C.: The Role of Diffusiophoresis in the Scavenging of Radioactive Particles from the Atmosphere, Q. J. Roy. Meteorol. Soc., 89, 43–61, doi:10.1002/qj.49708937903, 1963.

Gore, M. L., Aneja, V., Cooter, E. J., Dennis, R., and Pleim, J. E.: Ammonia Emissions in the U.S.: Assessing the role of bi-directional ammonia transport using the Community Multi-scale Air Quality (CMAQ) Model CEH, Centre of Ecology and Hydrology, Edinburgh, Scotland, UK, 2009.

Gupta, A., Tang, D., and McMurry, P. H., Growth of Monodisperse, Submicron Aerosol-Particles Exposed to SO₂, H₂O₂, and NH₃, J. Atmos. Chem., 20, 117–139, 1995.

Gyldenkerne, S., Skjøth, C. A., Hertel, O., and Ellermann, T.: A dynamical ammonia emission parameterization for use in air pollution models, J. Geophys. Res.-Atmos., 110, D07108, doi:10.1029/2004JD005459, 2005.

Hall, S. J., Matson, P. A., and Roth, P. M.: NO_x emissions from soil: Implications for air quality modeling in agricultural regions, Ann. Rev. Energy Environ., 21, 311–346, 1996.

Hampson, R. F. and Gavin, D.: Reaction rate and photochemical data for atmospheric chemistry – 1977, National Bureau of Standards, US Gov, 1978.

Harris, G. W., Klemp, D., and Zenker, T.: An Upper Limit on the HCl Near-Surface Mixing-Ratio Over the Atlantic Measured Using Tdlas, J. Atmos. Chem., 15, 327–332, 1992.

Harrison, R. M., Msibi, M. I., Kitto, A. M. N., and Yamulki, S.: Atmospheric Chemical-Transformations of Nitrogen-Compounds Measured in the North-Sea Experiment, September 1991, Atmos. Environ., 28, 1593–1599, 1994.

Harrison, R. M., Peak, J. D., and Collins, G. M.: Tropospheric cycle of nitrous acid, J. Geophys. Res.-Atmos., 101, 14429–14439, 1996.

Hertel, O.: Transformation and Deposition of Sulphur and Nitrogen Compounds in the Marine Boundary Layer, National Environmental Research Institute, Roskilde, Denmark, 215, 1–10, 1995.

Hertel, O. and Goodsite, M. E.: Urban Air Pollution Climate Through out the World in: Air Quality in Urban Environments, edited by: Hester, R. E. and Harrison, R. M., RSC Publishing, Cambridge, 1–22, 2009.

Hertel, O., Skjøth, C. A., Frohn, L. M., Vignati, E., Frydendall, J., de Leeuw, G., Schwarz, U., and Reis, S.: Assessment of the atmospheric nitrogen and sulphur inputs into the North Sea using a Lagrangian model, Phys. Chem. Earth, 27, 1507–1515, 2002.

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Hertel, O., Ambelas Skjøth, C., Brandt, J., Christensen, J. H., Frohn, L. M., and Frydendall, J.: Operational mapping of atmospheric nitrogen deposition to the Baltic Sea, *Atmos. Chem. Phys.*, 3, 2083–2099, doi:10.5194/acp-3-2083-2003, 2003.
- Hertel, O., Skjøth, C. A., Løfstrøm, P., Geels, C., Frohn, L. M., Ellermann, T., and Madsen, P. V.: Modelling Nitrogen Deposition on a Local Scale – A Review of the Current State of the Art, *Environ. Chem.*, 3, 317–337, 2006.
- Hertel, O., Hvidberg, M., Ketzel, M., Storm, L., and Stausgaard, L.: A proper choice of route significantly reduces air pollution exposure – A study on bicycle and bus trips in urban streets, *Sci. Total Environ.*, 389, 58–70, 2008.
- Hertel, O., Reis, S., Skjøth, C. A., Bleeker, A., Harrison, R. M., Cape, J. N., Fowler, D., Skiba, U., Simpson, D., Jickells, T., Baker, A. R., Kulmala, M., Gyldenkerne, S., Sørensen, L. L., and Erisman, J. W.: Nitrogen Processes in the Atmosphere in: *The European Nitrogen Assessment – Sources, Effects and Policy Perspectives*, edited by: Sutton, M. A., Howard, C. M., Erisman, J. W., Billen, G., Bleeker, A., Grennfelt, P., van Grinsven, H., and Grizzetti, B., Cambridge University Press, Cambridge, 177–207, 2011.
- Hicks, B. B., Baldocchi, D. D., Meyers, T. P., Hosker, R. P., and Matt, D. R.: A Preliminary Multiple Resistance Routine for Deriving Dry Deposition Velocities from Measured Quantities, *Water Air Soil Pollut.*, 36, 311–330, 1987.
- Hill, K. A., Shepson, P. B., Galbavy, E. S., Anastasio, C., Kourtev, P. S., Konopka, A., and Stirm, B. H.: Processing of atmospheric nitrogen by clouds above a forest environment, *J. Geophys. Res.-Atmos.*, 112, D11301, 2007.
- Horii, C. V., Munger, J. W., Wofsy, S. C., Zahniser, M., Nelson, D., and McManus, J. B.: Fluxes of nitrogen oxides over a temperate deciduous forest, *J. Geophys. Res.-Atmos.*, 109, D08305, doi:10.1029/2003JD004326, 2004.
- Horvath, L. and Sutton, M. A.: Long-term record of ammonia and ammonium concentrations at K-pusztá, Hungary, *Atmos. Environ.*, 32, 339–344, 1998.
- Horvath, L., Fagerli, H., and Sutton, M. A.: Long-term record (1981–2005) of ammonia and ammonium concentrations at K-pusztá Hungary and the effect of SO₂ emission change on measured and modelled concentrations, in: *Atmospheric Ammonia: Detecting Emission Changes and Environmental Impacts*, edited by: Sutton, M. A., Reis, S., and Baker, S. M. H., Springer Press, 181–186, 2009.
- Huntzicker, J. J., Cary, R. A., and Ling, C. S.: Neutralization of Sulfuric-Acid Aerosol by Ammonia, *Environ. Sci. Technol.*, 14, 819–824, 1980.

- Husted, S., Schjørring, J. K., Nielsen, K. H., Nemitz, E., and Sutton, M. A.: Stomatal compensation points for ammonia in oilseed rape plants under field conditions, *Agr. Forest Meteorol.*, 105, 371–383, 2000.
- Jenkin, M. E. and Clemitshaw, K. C.: Ozone and other secondary photochemical pollutants: chemical processes governing their formation in the planetary boundary layer, *Atmos. Environ.*, 34, 2499–2527, 2000.
- Joffre, S.: Parameterization and assessment of processes affecting the long-range transport of airborne pollutants over the sea, Finnish Meteorological Institute, Helsinki, Finland, 50, 1988.
- Johnson, M. T., Liss, P. S., Bell, T. G., Lesworth, T. J., Baker, A. R., Hind, A. J., Jickells, T. D., Biswas, K. F., Woodward, E. M. S., and Gibb, S. W.: Field observations of the ocean-atmosphere exchange of ammonia: Fundamental importance of temperature as revealed by a comparison of high and low latitudes, *Global Biogeochem. Cy.*, 22, GB1019, doi:10.1029/2007GB003039, 2008.
- Junge, E. and Ryan, T. G.: Study of the SO₂ Oxidation in Solution and Its Role in Atmospheric Chemistry, *Q. J. Roy. Meteorol. Soc.*, 84, 46–55, 1958.
- Kakosimos, K. E., Hertel, O., Ketzel, M., and Berkowicz, R.: Operational Street Pollution Model (OSPM) – a review of performed application and validation studies, and future prospects, *Environ. Chem.*, 7, 485–503, 2010.
- Kean, A. J., Littlejohn, D., Ban-Weiss, G. A., Harley, R. A., Kirchstetter, T. W., and Lunden, M. M.: Trends in on-road vehicle emissions of ammonia, *Atmos. Environ.*, 43, 1565–1570, 2009.
- Keene, W. C., Montag, J. A., Maben, J. R., Southwell, M., Leonard, J., Church, T. M., Moody, J. L., and Galloway, J. N.: Organic nitrogen in precipitation over Eastern North America, *Atmos. Environ.*, 36, 4529–4540, 2002.
- Krab, E. J., Cornelissen, J. H. C., Lang, S. I., and van Logtestijn, R. S. P.: Amino acid uptake among wide-ranging moss species may contribute to their strong position in higher-latitude ecosystems, *Plant Soil*, 304, 199–208, 2008.
- Kurtén, T., Loukonen, V., Vehkamäki, H., and Kulmala, M.: Amines are likely to enhance neutral and ion-induced sulfuric acid-water nucleation in the atmosphere more effectively than ammonia, *Atmos. Chem. Phys.*, 8, 4095–4103, doi:10.5194/acp-8-4095-2008, 2008.
- Lambrecht, U.: Legislation and future requirements for NO_x reduction Frankfurt, Germany, Paper in proceedings of the IQPC Conference, 2007.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Larsson, L., Ferm, M., Kasimir-Klemetsson, A., and Klemetsson, L.: Ammonia and nitrous oxide emissions from grass and alfalfa mulches, *Nutrient Cycl. Agroecosyst.*, 51, 41–46, 1998.
- Lee, D. S., Halliwell, C., Garland, J. A., Dollard, G. J., and Kingdon, R. D.: Exchange of ammonia at the sea surface – A preliminary study, *Atmos. Environ.*, 32, 431–439, 1998.
- Legrand, M., Preunkert, S., Wagenbach, D., and Fischer, H.: Seasonally resolved Alpine and Greenland ice core records of anthropogenic HCl emissions over the 20th century, *J. Geophys. Res.-Atmos.*, 107, 4139, doi:10.1029/2001JD001165, 2002.
- Leip, A., Achermann, B., Billen, G., Bleeker, A., Bouwman, A. F., de Vries, W., Dragosits, U., Döring, U., Fernall, D., Geupel, M., Herolstab, J., Johnes, P., le Gall, A. C., Monni, S., Neve-ceral, R., Orlandini, L., Prud'homme, M., Reuter, H. I., Simpson, D., Seufert, G., Spranger, T., Sutton, M. A., Van Aardenne, J., Voss, M., and Winiwarter, W.: Integrating nitrogen fluxes at the European scale in: *The European Nitrogen Assessment – Sources, Effects and Policy Perspectives*, edited by: Sutton, M. A., Howard, C. M., Erisman, J. W., Billen, G., Bleeker, A., Grennfelt, P., van Grinsven, H., and Grizzetti, B., Cambridge University Press, Cambridge, 345–376, 2011.
- Lipson, D. and Nasholm, T.: The unexpected versatility of plants: organic nitrogen use and availability in terrestrial ecosystems, *Oecologia*, 128, 305–316, 2001.
- Loubet, B., Milford, C., Sutton, M. A., and Cellier, P.: Investigation of the interaction between sources and sinks of atmospheric ammonia in an upland landscape using a simplified dispersion-exchange model, *J. Geophys. Res.-Atmos.*, 106, 24183–24195, 2001.
- Loubet, B., Asman, W. A. H., Theobald, M. R., Hertel, O., Tang, Y. S., Robin, P., Hassouna, M., Dammgén, U., Genermont, S., Cellier, P., and Sutton, M. A.: Ammonia deposition near hot spots: processes, models and monitoring methods in: *Atmospheric Ammonia: Detecting emission changes and environmental impacts*, edited by: Sutton, M. A., Reis, S., and Baker, S. M. H., Springer Press, 2009.
- Loubet, B., Genermont, S., Ferrara, R., Bedos, G., Decuq, G., Personne, E., Fanucci, O., Durand, B., Rana, G., and Cellier, P.: An inverse model to estimate ammonia emissions from fields, *Euro. J. Soil Sci.*, 61, 793–805, 2010.
- Loubet, B., Laville, P., Lehuger, S., Larmanou, E., Flechard, C., Mascher, N., Genermont, S., Roche, R., Ferrara, R. M., Stella, P., Personne, E., Durand, B., Decuq, C., Flura, D., Masson, S., Fanucci, O., Rampon, J. N., Siemens, J., Kindler, R., Gabrielle, B., Schruppf, M., and

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Cellier, P.: Carbon, nitrogen and Greenhouse gases budgets over a four years crop rotation in northern France, *Plant Soil*, 343, 109–137, 2011.
- Massad, R.-S., Nemitz, E., and Sutton, M. A.: Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere, *Atmos. Chem. Phys.*, 10, 10359–10386, doi:10.5194/acp-10-10359-2010, 2010a.
- Massad, R. S., Tuzet, A., Loubet, B., Perrier, A., and Cellier, P.: Model of stomatal ammonia compensation point (STAMP) in relation to the plant nitrogen and carbon metabolisms and environmental conditions, *Ecol. Model.*, 221, 479–494, 2010b.
- Mattsson, M., Herrmann, B., David, M., Loubet, B., Riedo, M., Theobald, M. R., Sutton, M. A., Bruhn, D., Neftel, A., and Schjoerring, J. K.: Temporal variability in bioassays of the stomatal ammonia compensation point in relation to plant and soil nitrogen parameters in intensively managed grassland, *Biogeosciences*, 6, 171–179, doi:10.5194/bg-6-171-2009, 2009.
- Mckay, H. A. C.: Atmospheric Oxidation of Sulphur Dioxide in Water Droplets in Presence of Ammonia, *Atmos. Environ.*, 5, 7–14, 1971.
- McMurry, P. H., Takano, H., and Anderson, G. R.: Study of the Ammonia (Gas) Sulfuric-Acid (Aerosol) Reaction-Rate, *Environ. Sci. Technol.*, 17, 347–352, 1983.
- Menut, L. and Bessagnet, B.: Atmospheric composition forecasting in Europe, *Ann. Geophys.*, 28, 61–74, doi:10.5194/angeo-28-61-2010, 2010.
- Milford, C., Theobald, M. R., Nemitz, E., Hargreaves, K. J., Horvath, L., Raso, J., Dämmgen, U., Neftel, A., Jones, S. K., Hensen, A., Loubet, B., Cellier, P., and Sutton, M. A.: Ammonia fluxes in relation to cutting and fertilization of an intensively managed grassland derived from an inter-comparison of gradient measurements, *Biogeosciences*, 6, 819–834, doi:10.5194/bg-6-819-2009, 2009.
- Misselbrook, T. H., Webb, J., Chadwick, D. R., Ellis, S., and Pain, B. F.: Gaseous emissions from outdoor concrete yards used by livestock, *Atmos. Environ.*, 35, 5331–5338, 2001.
- Monteith, J. and Unsworth, M.: *Principles of Environmental Physics*, Elsevier, New York, 2008.
- Muck, R. E. and Steenhuis, T. S.: Nitrogen Losses from Manure Storages, *Agr. Wastes*, 4, 41–54, 1982.
- Murphy, S. M., Sorooshian, A., Kroll, J. H., Ng, N. L., Chhabra, P., Tong, C., Surratt, J. D., Knipping, E., Flagan, R. C., and Seinfeld, J. H.: Secondary aerosol formation from atmospheric reactions of aliphatic amines, *Atmos. Chem. Phys.*, 7, 2313–2337, doi:10.5194/acp-7-2313-2007, 2007.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Neff, J. C., Holland, E. A., Dentener, F. J., McDowell, W. H., and Russell, K. M.: The origin, composition and rates of organic nitrogen deposition: A missing piece of the nitrogen cycle?, *Biogeochem.*, 57, 99–136, 2002.
- Neirynck, J., Kowalski, A. S., Carrara, A., and Ceulemans, R.: Driving forces for ammonia fluxes over mixed forest subjected to high deposition loads, *Atmos. Environ.*, 39, 5013–5024, 2005.
- Nemitz, E. and Sutton, M. A.: Gas-particle interactions above a Dutch heathland: III. Modelling the influence of the $\text{NH}_3\text{-HNO}_3\text{-NH}_4\text{NO}_3$ equilibrium on size-segregated particle fluxes, *Atmos. Chem. Phys.*, 4, 1025–1045, doi:10.5194/acp-4-1025-2004, 2004.
- Nemitz, E., Sutton, M. A., Gut, A., San Jose, R., Husted, S., and Schjørring, J. K.: Sources and sinks of ammonia within an oilseed rape canopy, *Agr. Forest Meteorol.*, 105, 385–404, 2000.
- Nemitz, E., Milford, C., and Sutton, M. A.: A two-layer canopy compensation point model for describing bi-directional biosphere-atmosphere exchange of ammonia, *Q. J. Roy. Meteorol. Soc.*, 127, 815–833, 2001.
- Nemitz, E., Sutton, M. A., Wyers, G. P., and Jongejan, P. A. C.: Gas-particle interactions above a Dutch heathland: I. Surface exchange fluxes of NH_3 , SO_2 , HNO_3 and HCl , *Atmos. Chem. Phys.*, 4, 989–1005, doi:10.5194/acp-4-989-2004, 2004a.
- Nemitz, E., Sutton, M. A., Wyers, G. P., Otjes, R. P., Mennen, M. G., van Putten, E. M., and Gallagher, M. W.: Gas-particle interactions above a Dutch heathland: II. Concentrations and surface exchange fluxes of atmospheric particles, *Atmos. Chem. Phys.*, 4, 1007–1024, doi:10.5194/acp-4-1007-2004, 2004b.
- Notholt, J., Hjorth, J., and Raes, F.: Formation of HNO_2 on aerosol surfaces during foggy periods in the presence of NO and NO_2 , *Atmos. Environ.*, 26A, 2111–2117, 1992.
- Olesen, J. E. and Sommer, S. G.: Modeling Effects of Wind-Speed and Surface Cover on Ammonia Volatilization from Stored Pig Slurry, *Atmos. Environ. Part A*, 27, 2567–2574, 1993.
- Oudendag, D. A. and Luesink, H. H.: The Manure Model: manure, minerals (N, P and K), ammonia emission, heavy metals and the use of fertiliser in Dutch agriculture, *Environ. Pollut.*, 102, 241–246, 1998.
- Paerl, H. W. and Whittall, D. R.: Anthropogenically-derived atmospheric nitrogen deposition, marine eutrophication and harmful algal bloom expansion: Is there a link?, *Ambio*, 28, 307–311, 1999.
- Palmgren, F., Berkowicz, R., Hertel, O., and Vignati, E.: Effects of reduction of NO_x on the NO_2 levels in urban streets, *Sci. Total Environ.*, 190, 409–415, 1996.

- Perrino, C., Catrambone, M., Di Bucchianico, A. D. M., and Allegrini, I., Gaseous ammonia in the urban area of Rome, Italy and its relationship with traffic emissions, *Atmos. Environ.*, 36, 5385–5394, 2002.
- Petaja, T., Sipila, M., Paasonen, P., Nieminen, T., Kurten, T., Ortega, I. K., Stratmann, F., Vehkamäki, H., Berndt, T., and Kulmala, M.: Experimental Observation of Strongly Bound Dimers of Sulfuric Acid: Application to Nucleation in the Atmosphere, *Phys. Rev. Lett.*, 106, 228302, doi:10.1103/PhysRevLett.106.228302, 2011.
- Petersen, S. O., Sommer, S. G., Aaes, O., and Sørengaard, K.: Ammonia losses from urine and dung of grazing cattle: effect of N intake, *Atmos. Environ.*, 32, 295–300, 1998.
- Petroff, A., Mailliat, A., Amielh, M., and Anselmetti, F.: Aerosol dry deposition on vegetative canopies, Part I: Review of present knowledge, *Atmos. Environ.*, 42, 3625–3653, 2008.
- Pilegaard, K., Skiba, U., Ambus, P., Beier, C., Brüggemann, N., Butterbach-Bahl, K., Dick, J., Dorsey, J., Duyzer, J., Gallagher, M., Gasche, R., Horvath, L., Kitzler, B., Leip, A., Pihlatie, M. K., Rosenkranz, P., Seufert, G., Vesala, T., Westrate, H., and Zechmeister-Boltenstern, S.: Factors controlling regional differences in forest soil emission of nitrogen oxides (NO and N₂O), *Biogeosciences*, 3, 651–661, doi:10.5194/bg-3-651-2006, 2006.
- Pinder, R. W., Adams, P. J., and Pandis, S. N.: Ammonia emission controls as a cost-effective strategy for reducing atmospheric particulate matter in the eastern United States, *Environ. Sci. Technol.*, 41, 380–386, 2007.
- Pinder, R. W., Davidson, E. A., Goodale, C. L., Greaver, T. L., Herrick, J. D., and Liu, L. L.: Climate change impacts of US reactive nitrogen, *Proc. Nat. Aca. Scie. USA*, 109, 7671–7675, 2012.
- Pio, C. A. and Harrison, R. M.: The Equilibrium of Ammonium-Chloride Aerosol with Gaseous Hydrochloric-Acid and Ammonia Under Tropospheric Conditions, *Atmos. Environ.*, 21, 1243–1246, 1987.
- Pouliot, G., Pierce, T., van der Gon, H. D., Schaap, M., Moran, M., and Nopmongkol, U.: Comparing emission inventories and model-ready emission datasets between Europe and North America for the AQMEII project, *Atmos. Environ.*, 53, 4–14, 2012.
- Pruppacher, H. R. and Jaenicke, R.: The Processing of Water-Vapor and Aerosols by Atmospheric Clouds, *A Global Estimate*, *Atmos. Res.*, 38, 283–295, 1995.
- Pryor, S. C. and Sorensen, L. L.: Nitric acid-sea salt reactions: Implications for nitrogen deposition to water surfaces, *J. Appl. Meteorol.*, 39, 725–731, 2000.

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Pryor, S. C., Barthelmie, R. J., Sørensen, L. L., Larsen, S. E., Sempreviva, A. M., Grönholm, T., Rannik, Ü., Kulmala, M., and Vesala, T.: Upward fluxes of particles over forests: when, where, why?, *Tellus B*, 60, 372–380, 2008a.
- Pryor, S. C., Gallagher, M., Sievering, H., Larsen, S. E., Barthelmie, R. J., Birsan, F., Nemitz, E., Rinne, J., Kulmala, M., Grünholm, T., Taipale, R., and Vesala, T.: A review of measurement and modelling results of particle atmosphere↔surface exchange, *Tellus B*, 60, 42–75, 2008b.
- Qualls, R. G. and Haines, B. L.: Biodegradability of Dissolved Organic-Matter in Forest Throughfall, Soil Solution, and Stream Water, *Soil Sci. Soc. Am. J.*, 56, 578–586, 1992.
- Quinn, P. K., Charlson, R. J., and Bates, T. S.: Simultaneous Observations of Ammonia in the Atmosphere and Ocean, *Nature*, 335, 336–338, 1988.
- Raes, F., Van Dingenen, R., Vignati, E., Wilson, J., Putaud, J. P., Seinfeld, J. H., and Adams, P.: Formation and cycling of aerosols in the global troposphere, *Atmos. Environ.*, 34, 4215–4240, 2000.
- Reche, C., Viana, M., Pandolfi, M., Alastuey, A. S., Moreno, T., Amato, F., Ripoll, A., and Querol, X.: Urban NH₃ levels and sources in a Mediterranean environment, *Atmos. Environ.*, 57, 153–164, 2012.
- Reis, S., Pinder, R. W., Zhang, M., Lijie, G., and Sutton, M. A.: Reactive nitrogen in atmospheric emission inventories, *Atmos. Chem. Phys.*, 9, 7657–7677, doi:10.5194/acp-9-7657-2009, 2009.
- Riddick, S. N., Dragosits, U., Blackall, T. D., Daunt, F., Wanless, S., and Sutton, M. A.: The global distribution of ammonia emissions from seabird colonies, *Atmos. Environ.*, 55, 319–327, 2012.
- Riedo, M., Milford, C., Schmid, M., and Sutton, M. A.: Coupling soil-plant-atmosphere exchange of ammonia with ecosystem functioning in grasslands, *Ecol. Model.*, 158, 83–110, 2002.
- Rierner, N., Vogel, H., Vogel, B., Anttila, T., Kiendler-Scharr, A., and Mentel, T. F.: Relative importance of organic coatings for the heterogeneous hydrolysis of N₂O₅ during summer in Europe, *J. Geophys. Res.-Atmos.*, 114, D17307, doi:10.1029/2008JD011369, 2009.
- Ryden, J. C., Whitehead, D. C., Lockyer, D. R., Thompson, R. B., Skinner, J. H., and Garwood, E. A.: Ammonia Emission from Grassland and Livestock Production Systems in the UK, *Environ. Pollut.*, 48, 173–184, 1987.
- Schade, G. W. and Crutzen, P. J.: Emission of Aliphatic-Amines from Animal Husbandry and Their Reactions - Potential Source of N₂O and Hcn, *J. Atmos. Chem.*, 22, 319–346, 1995.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏮

⏭

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Schjørring, J. K., Husted, S., and Poulsen, M. M.: Soil-plant-atmosphere ammonia exchange associated with *Calluna vulgaris* and *Deschampsia flexuosa*, *Atmos. Environ.*, 32, 507–512, 1998.

Schwarz, U., Wickert, B., Obermeier, A., and Friedrich, R.: Generation of Atmospheric Emission Inventories in Europe with High Spatial and Temporal Resolution, edited by: Borrell, P. M. and Borrell, P., WITpress, Southampton, Southampton SO40 7AA, UK, 2000.

Seedorf, J., Hartung, J., Schroder, M., Linkert, K. H., Pedersen, S., Takai, H., Johnsen, J. O., Metz, J. H. M., Groot Koerkamp, P. W. G., and Uenk, G. H.: Temperature and Moisture Conditions in Livestock Buildings in Northern Europe, *J. Agr. Eng. Res.*, 70, 49–57, 1998.

Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, John Wiley & Sons Inc., New York, 1203 pp., 2006.

Simpson, D., Winiwarter, W., Borjesson, G., Cinnerby, S., Ferreira, A., Guenther, A., Hewitt, C. N., Janson, R., Khalil, M. A. K., Owen, S., Pierce, T. E., Puxbaum, H., Shearer, M., Skiba, U., Steinbrecher, R., Tarrason, L., and Oquist, M. G.: Inventorying emissions from nature in Europe, *J. Geophys. Res.-Atmos.*, 104, 8113–8152, 1999.

Simpson, D., Ashmore, M. R., Emberson, L., and Tuovinen, J. P.: A comparison of two different approaches for mapping potential ozone damage to vegetation. A model study, *Environ. Pollut.*, 146, 715–725, 2007.

Simpson, D., Aas, W., Bartnicki, J., Berge, H., Bleeker, A., Cuvelier, C., Dentener, F., Dore, T., Erismann, J. W., Fagerli, H., Flechard, C., Hertel, O., Van Jaarsveld, H., Jenkin, M. E., Schaap, M., Smeena, V. S., Thunis, P., Vautard, R., and Vieno, M.: Chapter 14: Atmospheric transport and deposition of reactive nitrogen in Europe in: *The European Nitrogen Assessment – Sources, Effects and Policy Perspectives*, edited by: Sutton, M., Howard, C. M., Erismann, J. W., Billen, G., Bleeker, A., Grennfelt, P., van Grinsven, H., and Grizzetti, B., Cambridge University Press, Cambridge, UK, 298–316, 2011.

Simpson, D., Benedictow, A., Berge, H., Bergström, R., Emberson, L. D., Fagerli, H., Hayman, G. D., Gauss, M., Jonson, J. E., Jenkin, M. E., Nyíri, A., Richter, C., Semeena, V. S., Tsyro, S., Tuovinen, J.-P., Valdebenito, Á., and Wind, P.: The EMEP MSC-W chemical transport model – Part 1: Model description, *Atmos. Chem. Phys. Discuss.*, 12, 3781–3874, doi:10.5194/acpd-12-3781-2012, 2012.

Sintermann, J., Neftel, A., Ammann, C., Häni, C., Hensen, A., Loubet, B., and Flechard, C. R.: Are ammonia emissions from field-applied slurry substantially over-estimated in European emission inventories?, *Biogeosciences*, 9, 1611–1632, doi:10.5194/bg-9-1611-2012, 2012.

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Skiba, U., Fowler, D., and Smith, K. A.: Nitric oxide emissions from agricultural soils in temperate and tropical climates: sources, controls and mitigation options, *Nutrient Cycl. Agroecosyst.*, 48, 139–153, 1997.

Skiba, U., Drewer, J., Tang, Y. S., van Dijk, N., Helfter, C., Nemitz, E., Famulari, D., Cape, J. N., Jones, S. K., Twigg, M., Pihlatie, M., Vesala, T., Larsen, K. S., Carter, M. S., Ambus, P., Ibrom, A., Beier, C., Hensen, A., Frumau, A., Erismann, J. W., Bruggemann, N., Gasche, R., Butterbach-Bahl, K., Neftel, A., Spirig, C., Horvath, L., Freibauer, A., Cellier, P., Laville, P., Loubet, B., Magliulo, E., Bertolini, T., Seufert, G., Andersson, M., Manca, G., Laurila, T., Aurela, M., Lohila, A., Zechmeister-Boltenstern, S., Kitzler, B., Schauffler, G., Siemens, J., Kindler, R., Flechard, C., and Sutton, M. A.: Biosphere-atmosphere exchange of reactive nitrogen and greenhouse gases at the NitroEurope core flux measurement sites: Measurement strategy and first data sets, *Agr. Ecosyst. Environ.*, 133, 139–149, 2009.

Skjøth, C. A., Hertel, O., and Ellermann, T.: Use of the ACDEP trajectory model in the Danish nation-wide Background Monitoring Programme, *Phys. Chem. Earth*, 27, 1469–1477, 2002.

Skjøth, C. A., Hertel, O., Gyldenkerne, S., and Ellermann, T.: Implementing a dynamical ammonia emission parameterization in the large-scale air pollution model ACDEP, *J. Geophys. Res.-Atmos.*, 109, D06306, doi:10.1029/2003JD003895, doi:10.1029/2003JD003895, 2004.

Skjøth, C. A., Ellermann, T., Hertel, O., Gyldenkerne, S., and Mikkelsen, M. H.: Footprints on ammonia concentrations from environmental regulations, *J. Air Waste Managem. Assoc.*, 58, 1158–1165, 2008.

Skjøth, C. A., Geels, C., Berge, H., Gyldenkerne, S., Fagerli, H., Ellermann, T., Frohn, L. M., Christensen, J., Hansen, K. M., Hansen, K., and Hertel, O.: Spatial and temporal variations in ammonia emissions – a freely accessible model code for Europe, *Atmos. Chem. Phys.*, 11, 5221–236, doi:10.5194/acp-11-5221-2011, 2011.

Slinn, W. G. N.: Predictions for Particle Deposition to Vegetative Canopies, *Atmos. Environ.*, 16, 1785–1794, 1982.

Smith, E., Gordon, R., Bourque, C., Campbell, A., Genermont, S., Rochette, P., and Mkhabela, M.: Simulated management effects on ammonia emissions from field applied manure, *J. Environ. Manage.*, 90, 2531–2536, 2009.

Smith, R. I., Fowler, D., Sutton, M. A., Flechard, C., and Coyle, M.: Regional estimation of pollutant gas dry deposition in the UK: model description, sensitivity analyses and outputs, *Atmos. Environ.*, 34, 3757–3777, 2000.

- Sommer, S. G., Østergård, H. S., Løfstrøm, P., Andersen, H. V., and Jensen, L. S.: Validation of model calculation of ammonia deposition in the neighbourhood of a poultry farm using measured NH_3 concentrations and N deposition, *Atmos. Environ.*, 43, 915–920, 2009.
- Sørensen, L. L., Hertel, O., Skjøth, C. A., Lund, M., and Pedersen, B.: Fluxes of ammonia in the coastal marine boundary layer, *Atmos. Environ.*, 37, S167–S177, 2003.
- Sorensen, L. L., Pryor, S. C., de Leeuw, G., and Schulz, M.: Flux divergence of nitric acid in the marine atmospheric surface layer, *J. Geophys. Res.-Atmos.*, 110, D15306, doi:10.1029/2004JD005403, 2005.
- Sorteberg, A. and Hov, Ø.: Two parametrizations of the dry deposition exchange for SO_2 and NH_3 in a numerical model, *Atmos. Environ.*, 30, 1823–1840, 1996.
- Spokes, L., Jickells, T., Weston, K., Gustafsson, B. G., Johnsson, M., Liljebladh, B., Conley, D., Skjøth, C. A., Brandt, J., Carstensen, J., Christiansen, T., Frohn, L., Geernaert, G., Hertel, O., Jensen, B., Lundsgaard, C., Markager, S., Martinsen, W., Møller, B., Pedersen, B., Sauerberg, K., Sorensen, L. L., Hasager, C. C., Sempreviva, A. M., Pryor, S. C., Lund, S. W., Larsen, S., Tjernstrøm, M., Svensson, G., and Zagar, M.: MEAD: An interdisciplinary study of the marine effects of atmospheric deposition in the Kattegat, *Environ. Pollut.*, 140, 453–462, 2006.
- Stelson, A. W. and Seinfeld, J. H.: Thermodynamic Prediction of the Water Activity, NH_4NO_3 Dissociation-Constant, Density and Refractive-Index for the NH_4NO_3 -(NH_4) 2SO_4 - H_2O System at 25-Degrees-C, *Atmos. Environ.*, 16, 2507–2514, 1982.
- Stelson, A. W., Friedlander, S. K., and Seinfeld, J. H.: Note on the Equilibrium Relationship Between Ammonia and Nitric-Acid and Particulate Ammonium-Nitrate, *Atmos. Environ.*, 13, 369–371, 1979.
- Stemmler, K., Ammann, M., Donders, C., Kleffmann, J., and George, C.: Photosensitized reduction of nitrogen dioxide on humic acid as a source of nitrous acid, *Nature*, 440, 195–198, 2006.
- Stohl, A., Williams, E., Wotawa, G., and KrompKolb, H.: A European inventory of soil nitric oxide emissions and the effect of these emissions on the photochemical formation of ozone, *Atmos. Environ.*, 30, 3741–3755, 1996.
- Su, H., Cheng, Y. F., Oswald, R., Behrendt, T., Trebs, I., Meixner, F. X., Andreae, M. O., Cheng, P., Zhang, Y., and Poschl, U.: Soil Nitrite as a Source of Atmospheric HONO and OH Radicals, *Science*, 333, 1616–1618, 2011.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Sutton, M. A. and Fowler, D.: A model for inferring bi-directional fluxes of ammonia over plant canopies, Paper in the proceedings of the WMO conference on the measurement and modelling of atmospheric composition changes including pollutant transport, Sofia, Bulgaria, 1993.
- 5 Sutton, M. A., Fowler, D., Moncrieff, J. B., and Storetonwest, R. L.: The Exchange of Atmospheric Ammonia with Vegetated Surfaces .2. Fertilized Vegetation, Q. J. Roy. Meteorol. Soc., 119, 1047–1070, 1993.
- Sutton, M. A., Schjorring, J. K., and Wyers, G. P.: Plant Atmosphere Exchange of Ammonia, Philos. Trans. Roy. Soc. London, 351, 261–276, 1995.
- 10 Sutton, M. A., Burkhardt, J. K., Guerin, D., Nemitz, E., and Fowler, D.: Development of resistance models to describe measurements of bi-directional ammonia surface-atmosphere exchange, Atmos. Environ., 32, 473–480, 1998.
- Sutton, M. A., Dragosits, U., Tang, Y. S., and Fowler, D.: Ammonia emissions from non-agricultural sources in the UK, Atmos. Environ., 34, 855–869, 2000a.
- 15 Sutton, M. A., Nemitz, E., Fowler, D., Wyers, G. P., Otjes, R. P., Schjorring, J. K., Husted, S., Nielsen, K. H., San Jose, R., Moreno, J., Gallagher, M. W., and Gut, A.: Fluxes of ammonia over oilseed rape – Overview of the EXAMINE experiment, Agr. Forest Meteorol., 105, 327–349, 2000b.
- Sutton, M. A., Asman, W. A. H., Ellermann, T., Van Jaarsveld, J. A., Acker, K., Aneja, V., Duyzer, J., Horvath, L., Paramonov, S., Mitosinkova, M., Tang, Y. S., Achermann, B., Gauger, T., Bartnik, J., Neftel, A., and Erisman, J. W.: Establishing the link between ammonia emission control and measurements of reduced nitrogen concentrations and deposition, Environ. Monitor. Assess., 82, 149–185, 2003.
- 20 Sutton, M. A., Nemitz, E., Erisman, J. W., Beier, C., Bahl, K. B., Cellier, P., de Vries, W., Cotrufo, F., Skiba, U., Di Marco, C., Jones, S., Laville, P., Soussana, J. F., Loubet, B., Twigg, M., Famulari, D., Whitehead, J., Gallagher, M. W., Neftel, A., Flechard, C. R., Herrmann, B., Calanca, P. L., Schjoerring, J. K., Daemmgen, U., Horvath, L., Tang, Y. S., Emmett, B. A., Tietema, A., Penuelas, J., Kesik, M., Brueggemann, N., Pilegaard, K., Vesala, T., Campbell, C. L., Olesen, J. E., Dragosits, U., Theobald, M. R., Levy, P., Mobbs, D. C., Milne, R., Viovy, N., Vuichard, N., Smith, J. U., Smith, P., Bergamaschi, P., Fowler, D., and Reis, S.: Challenges in quantifying biosphere-atmosphere exchange of nitrogen species, Environ. Pollut., 150, 125–139, 2007.
- 25
- 30

- Sutton, M. A., Erisman, J. W., Dentener, F., and Moller, D.: Ammonia in the environment: From ancient times to the present, *Environ. Pollut.*, 156, 583–604, 2008.
- Sutton, M. A., Nemitz, E., Milford, C., Campbell, C., Erisman, J. W., Hensen, A., Cellier, P., David, M., Loubet, B., Personne, E., Schjoerring, J. K., Mattsson, M., Dorsey, J. R., Gallagher, M. W., Horvath, L., Weidinger, T., Meszaros, R., Dämmgen, U., Neftel, A., Herrmann, B., Lehman, B. E., Flechard, C., and Burkhardt, J.: Dynamics of ammonia exchange with cut grassland: synthesis of results and conclusions of the GRAMINAE Integrated Experiment, *Biogeosciences*, 6, 2907–2934, doi:10.5194/bg-6-2907-2009, 2009a.
- Sutton, M. A., Nemitz, E., Theobald, M. R., Milford, C., Dorsey, J. R., Gallagher, M. W., Hensen, A., Jongejan, P. A. C., Erisman, J. W., Mattsson, M., Schjoerring, J. K., Cellier, P., Loubet, B., Roche, R., Neftel, A., Hermann, B., Jones, S. K., Lehman, B. E., Horvath, L., Weidinger, T., Rajkai, K., Burkhardt, J., Löpmeier, F. J., and Daemmgen, U.: Dynamics of ammonia exchange with cut grassland: strategy and implementation of the GRAMINAE Integrated Experiment, *Biogeosciences*, 6, 309–331, doi:10.5194/bg-6-309-2009, 2009b.
- Sutton, M. A., Reis, S., and Baker, S. M. H.: Atmospheric Ammonia – Detecting emission changes and environmental impacts – Results from an Expert Workshop under the Convention on Long-range Transboundary Air Pollution, Springer Publishers, 464 pp., 2009c.
- Sutton, M. A., Oenema, O., Erisman, J. W., Leip, A., van Grinsven, H., and Winiwarter, W.: Too much of a good thing, *Nature*, 472, 159–161, 2011.
- Tang, Y. S., Simmons, I., van Dijk, N., Di Marco, C., Nemitz, E., Demmgen, U., Gilke, K., Djuricic, V., Vidic, S., Gliha, Z., Borovecki, D., Mitosinkova, M., Hanssen, J. E., Uggerud, T. H., Sanz, M. J., Sanz, P., Chorda, J. V., Flechard, C. R., Fauvel, Y., Ferm, M., Perrino, C., and Sutton, M. A.: European scale application of atmospheric reactive nitrogen measurements in a low-cost approach to infer dry deposition fluxes, *Agriculture, Ecosyst. Environ.*, 133, 183–195, 2009.
- The European Nitrogen Assessment: Sources, Effects and Policy Perspectives, Cambridge University Press, Cambridge, 612 pp., 2011.
- Theobald, M. R., Crittenden, P. D., Hunt, A. P., Tang, Y. S., Dragosits, U., and Sutton, M. A.: Ammonia emissions from a Cape fur seal colony, Cape Cross, Namibia, *Geophys. Res. Lett.*, 33, L03812, doi:10.1029/2005GL024384, 2006.
- Theobald, M. R., Bealey, W. J., Tang, Y. S., Vallejo, A., and Sutton, M. A.: A simple model for screening the local impacts of atmospheric ammonia, *Sci. Total Environ.*, 407, 6024–6033, 2009.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Thoene, B., Schroder, P., Papen, H., Egger, A., and Rennenberg, H.: Absorption of Atmospheric NO₂ by Spruce (*Picea-Abies* L Karst) Trees .1. NO₂ Influx and Its Correlation with Nitrate Reduction, *New Phytologist*, 117, 575–585, 1991.

Tørseth, K., Aas, W., Breivik, K., Fjæraa, A. M., Fiebig, M., Hjellbrekke, A. G., Lund Myhre, C., Solberg, S., and Yttri, K. E.: Introduction to the European Monitoring and Evaluation Programme (EMEP) and observed atmospheric composition change during 1972–2009, *Atmos. Chem. Phys.*, 12, 5447–5481, doi:10.5194/acp-12-5447-2012, 2012.

Trebs, I., Lara, L. L., Zeri, L. M. M., Gatti, L. V., Artaxo, P., Dlugi, R., Slanina, J., Andreae, M. O., and Meixner, F. X.: Dry and wet deposition of inorganic nitrogen compounds to a tropical pasture site (Rondônia, Brazil), *Atmos. Chem. Phys.*, 6, 447–469, doi:10.5194/acp-6-447-2006, 2006.

Tunnicliffe, W. S., Burge, P. S., and Ayres, J. G.: Effect of Domestic Concentrations of Nitrogen-Dioxide on Airway Responses to Inhaled Allergen in Asthmatic-Patients, *Lancet*, 344, 1733–1736, 1994.

Turnipseed, A. A., Huey, L. G., Nemitz, E., Stickel, R., Higgs, J., Tanner, D. J., Slusher, D. L., Sparks, J. P., Flocke, F., and Guenther, A.: Eddy covariance fluxes of peroxyacetyl nitrates (PANs) and NO_y to a coniferous forest, *J. Geophys. Res.-Atmos.*, 111, D09304, doi:10.1029/2005JD006631, 2006.

van Glasow, R. and Crutzen, P.: Tropospheric halogen chemistry in: *Treatise on Geochemistry*, edited by: Holland, D. H. and Turekian, K. K., Elsevier-Pergamon, Oxford, UK, 1–67, 2007.

van Vuuren, D. P., Bouwman, L. F., Smith, S. J., and Dentener, F.: Global projections for anthropogenic reactive nitrogen emissions to the atmosphere: an assessment of scenarios in the scientific literature, *Curr. Opinion Environ. Sustain.*, 3, 359–369, 2011.

Vestreng, V., Ntziachristos, L., Semb, A., Reis, S., Isaksen, I. S. A., and Tarrasón, L.: Evolution of NO_x emissions in Europe with focus on road transport control measures, *Atmos. Chem. Phys.*, 9, 1503–1520, doi:10.5194/acp-9-1503-2009, 2009.

von Kuhlmann, R. and Lawrence, M. G.: The impact of ice uptake of nitric acid on atmospheric chemistry, *Atmos. Chem. Phys.*, 6, 225–235, doi:10.5194/acp-6-225-2006, 2006.

Wall, S. M., John, W., and Ondo, J. L.: Measurements of aerosol size distributions for nitrate and major ionic species, *Atmos. Environ.*, 22, 1649–1656, 1988.

Wayne, R. P., Barnes, I., Biggs, P., Burrows, J. P., Canosamas, C. E., Hjorth, J., Lebras, G., Moortgat, G. K., Perner, D., Poulet, G., Restelli, G., and Sidebottom, H.: The Nitrate Radical – Physics, Chemistry, and the Atmosphere, *Atmos. Environ.*, 25, 1–203, 1991.

BGD

9, 9349–9423, 2012

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Webb, J., Anthony, S. G., Brown, L., Lyons-Visser, H., Ross, C., Cottrill, B., Johnson, P., and Scholefield, D.: The impact of increasing the length of the cattle grazing season on emissions of ammonia and nitrous oxide and on nitrate leaching in England and Wales, *Agr. Ecosyst. Environ.*, 105, 307–321, 2005.
- 5 Wells, M., Bower, K. N., Choularton, T. W., Cape, J. N., Sutton, M. A., Storetonwest, R. L., Fowler, D., Wiedensohler, A., Hansson, H. C., Svenningsson, B., Swietlicki, E., Wendisch, M., Jones, B., Dollard, G., Acker, K., Wieprecht, W., Preiss, M., Arends, B. G., Pahl, S., Berner, A., Kruisz, C., Laj, P., Facchini, M. C., and Fuzzi, S.: The reduced nitrogen budget of an orographic cloud, *Atmos. Environ.*, 31, 2599–2614, 1997.
- 10 Wichink Kruit, R. J., Schaap, M., Sauter, F. J., van Zanten, M. C., and van Pul, W. A. J.: Modelling the distribution of ammonia across Europe including bi-directional surface-atmosphere exchange, *Biogeosciences Discuss.*, 9, 4877–4918, doi:10.5194/bgd-9-4877-2012, 2012.
- Wickert, B., Heidekker, A., and Friedrich, R.: Calculations of emissions in Europe with CAREAIR, edited by: Migley, P. M., Reuther, M., and Williams, M., Springer-Verlag, Berlin, Heidelberg, New York, 2001.
- 15 Wilson, L. J., Bacon, P. J., Bull, J., Dragosits, U., Blackall, T. D., Dunn, T. E., Hamer, K. C., Sutton, M. A., and Wanless, S.: Modelling the spatial distribution of ammonia emissions from seabirds in the UK, *Environ. Pollut.*, 131, 173–185, 2004.
- Wolff, V., Trebs, I., Foken, T., and Meixner, F. X.: Exchange of reactive nitrogen compounds: concentrations and fluxes of total ammonium and total nitrate above a spruce canopy, *Biogeosciences*, 7, 1729–1744, doi:10.5194/bg-7-1729-2010, 2010.
- 20 Wu, Z. Y., Wang, X. M., Turnipseed, A. A., Chen, F., Zhang, L. M., Guenther, A. B., Karl, T., Huey, L. G., Niyogi, D., Xia, B. C., and Alapaty, K.: Evaluation and improvements of two community models in simulating dry deposition velocities for peroxyacetyl nitrate (PAN) over a coniferous forest, *J. Geophys. Res.-Atmos.*, 117, D04310, doi:10.1029/2011JD016751, 2012.
- 25 Xu, L. and Penner, J. E.: Global simulations of nitrate and ammonium aerosols and their radiative effects, *Atmos. Chem. Phys. Discuss.*, 12, 10115–10179, doi:10.5194/acpd-12-10115-2012, 2012.
- Yeatman, S. G., Spokes, L. J., and Jickells, T. D.: Comparisons of coarse-mode aerosol nitrate and ammonium at two polluted coastal sites, *Atmos. Environ.*, 35, 1321–1335, 2001.
- 30 Yokelson, R. J., Urbanski, S. P., Atlas, E. L., Toohey, D. W., Alvarado, E. C., Crounse, J. D., Wennberg, P. O., Fisher, M. E., Wold, C. E., Campos, T. L., Adachi, K., Buseck, P. R., and

- Hao, W. M.: Emissions from forest fires near Mexico City, *Atmos. Chem. Phys.*, 7, 5569–5584, doi:10.5194/acp-7-5569-2007, 2007.
- Zhang, L., Wright, L. P., and Asman, W. A. H.: Bi-directional air-surface exchange of atmospheric ammonia: A review of measurements and a development of a big-leaf model for applications in regional-scale air-quality models, *J. Geophys. Res.-Atmos.*, 115, D20310, doi:10.1029/2009JD013589, 2010.
- Zhang, L. M., Gong, S. L., Padro, J., and Barrie, L.: A size-segregated particle dry deposition scheme for an atmospheric aerosol module, *Atmos. Environ.*, 35, 549–560, 2001.
- Zhang, Q. and Anastasio, C.: Chemistry of fog waters in California's Central Valley – Part 3: concentrations and speciation of organic and inorganic nitrogen, *Atmos. Environ.*, 35, 5629–5643, 2001.
- Zhang, R. H., Day, D. L., Christianson, L. L., and Jepson, W. P.: A Computer-Model for Predicting Ammonia Release Rates from Swine Manure Pits, *J. Agr. Eng. Res.*, 58, 223–229, 1994.
- Zhang, Y., Song, L., Liu, X. J., Li, W. Q., Lu, S. H., Zheng, L. X., Bai, Z. C., Cai, G. Y., and Zhang, F. S.: Atmospheric organic nitrogen deposition in China (vol 46C, pg 195, 2011), *Atmos. Environ.*, 49, 195–204, doi:10.1016/j.atmosenv.2011.09.080, 2012.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Governing processes for reactive nitrogen compounds

O. Hertel et al.

Table 1. Comparison of European emissions of N_r in Tg, using the nitrogen content of both NH_3 and NO_x to analyse the overall trend development of N_r release into the atmosphere. Emission figures based on the EDGAR* emission inventory, v4.2.

	1985		1990		1995		2000		2005	
	Tg	Share (%)	Tg	Share (%)	Tg	Share (%)	Tg	Share (%)	Tg	Share (%)
Agriculture	5.98	55.3 %	5.68	54.2 %	4.80	52.8 %	4.76	55.3 %	4.65	55.5 %
Road transport	1.73	16.0 %	2.09	20.0 %	1.85	20.3 %	1.51	17.6 %	1.34	15.9 %
Other mobile sources	0.36	3.3 %	0.26	2.5 %	0.24	2.6 %	0.24	2.8 %	0.26	3.1 %
Power generation	1.43	13.2 %	1.38	13.2 %	1.31	14.4 %	1.18	13.8 %	1.27	15.2 %
Industrial production	0.87	8.0 %	0.65	6.2 %	0.53	5.9 %	0.51	5.9 %	0.48	5.8 %
Other	0.44	4.1 %	0.41	3.9 %	0.36	4.0 %	0.40	4.6 %	0.38	4.5 %
Total	10.81		10.47		9.08		8.60		8.38	
			–3 %		–16 %		–20 %		–23 %	

* Source: European Commission, Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL). Emission Database for Global Atmospheric Research (EDGAR), release version 4.2.
<http://edgar.jrc.ec.europa.eu>, 2011.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Governing processes for reactive nitrogen compounds

O. Hertel et al.




Fig. 1. A sketch which illustrates the various pathways of reactive nitrogen compounds in the atmosphere. The left side of the figure illustrates the atmospheric pathways of NH_x (gas phase ammonia (NH₃) and aerosol phase ammonium (NH₄⁺)), and the right side of the figure illustrates the pathway of nitrogen oxides (NO_y) compounds in the atmosphere (NO_x = NO + NO₂ and reaction products). The sketch is a modified version of the figure in Hertel et al. (2006).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 2. The indoor temperature as function of outdoor ambient temperature for isolated stables (black) and open barns (pink). Isolated stables have a less steep increase in indoor temperatures because the ventilation rate (red) in these stable systems depends on outside temperatures.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Simulations of ammonia emission for open and isolated stables and manure storage using hourly meteorological input of temperature and wind speed. All times series were created by using meteorological data for the year the year 2007 and the emission model available at <http://www.atmos-chem-phys.net/11/5221/2011/acp-11-5221-2011.html>.

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Spatial distributions of annual NH_3 emissions in Europe for the year 2000 based on: (left) EMEP at $50 \text{ km} \times 50 \text{ km}$ to the right) EDGAR at $0.1^\circ \times 0.1^\circ$.

Governing processes for reactive nitrogen compounds

O. Hertel et al.




Fig. 5. Development of emissions of nitrogen oxides (left) and ammonia (right) in three world regions (Europe, USA, PR China) between 1970 and 2008 based on the Emissions Database for Global Atmospheric Research (EDGAR) version 4.2 (Unit: Tg). For a more detailed analysis of N_x emission trends, see Reis et al. (2009). Source: European Commission, Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL). Emission Database for Global Atmospheric Research (EDGAR), release version 4.2. <http://edgar.jrc.ec.europa.eu>, 2011

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Governing processes for reactive nitrogen compounds

O. Hertel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 6. Spatial distributions of annual NO_x emissions in Europe for the year 2000 based on: (left) EMEP at 50 km × 50 km to the right) EDGAR at 0.1 × 0.1°.

Governing processes for reactive nitrogen compounds

O. Hertel et al.




Fig. 7. Illustration of the interaction between the various nitrogen oxide (NO_y) compounds in the tropospheric boundary layer. The symbol Δ represents energy leading to thermal degradation, $h\nu$ solar radiation leading to photo dissociation and RH a hydrocarbon reacting with the species in question. PPN is a notation for other peroxy nitrates than PAN. Source (Derwent and Hertel, 1998).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 8. Farm scale NH₃ emission and deposition, illustrating the rapid fall-off in deposition levels with distance from source (adapted from Fowler et al., 1998 and modified from version presented in Simpson et al., 2011).

Governing processes for reactive nitrogen compounds

O. Hertel et al.




Fig. 9. Comparison of modelled and observed annual wet deposition of to the left NH_x and to the right NO_3^- (HNO_3 and aerosol phase NO_3^-). Data are for 2001 in the EMEP model with observations. The bullets depict observations with the same colour bas as the modelled field. Measured annual deposition is calculated by using the measured precipitation amount and the NO_3^- and NH_4^+ concentration in precipitation. Source: Simpson et al. (2011).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion